

Locally Anisotropic Gravity and Strings

Sergiu I. Vacaru

Institute of Applied Physics, Academy of Sciences of Moldova

5 Academy str., Chişinău 277028, Republic of Moldova

(February 7, 2008)

Abstract

An introduction into the theory of locally anisotropic spaces (modelled as vector bundles provided with compatible nonlinear and distinguished linear connections and metric structures and containing as particular cases different types of Kaluza–Klein and/or extensions of Lagrange and Finsler spaces) is presented. The conditions for consistent propagation of closed strings in locally anisotropic background spaces are analyzed. The connection between the conformal invariance, the vanishing of the renormalization group β -function of the generalized σ -model and field equations of locally anisotropic gravity is studied in detail.

© S.I. Vacaru

PACS numbers: 11.25.W, 11.25.Db, 11.10.Ln, 11.10.Gh

Typeset using REVTeX

I. INTRODUCTION

The relationship between two dimensional σ -models and strings has been considered by several authors [1-5] in order to discuss the effective low energy field equations for the massless models of strings. In this paper we shall investigate some of the problems associated with the theory of locally anisotropic strings being a natural generalization to locally anisotropic (la) backgrounds (we shall write in brief la-backgrounds, la-spaces and la-geometry) of the Polyakov's covariant functional-integral approach to string theory [6]. Our aim is to show that a corresponding low-energy string dynamics contains the motion equations for field equations on la-spaces.

The first geometric models of la-spaces were formulated by P. Finsler [7], studied in detail and generalized by A. Berwald [8] and E. Cartan [9]. The geometry of Lagrange and Finsler spaces, its extensions and possible applications in physics (locally anisotropic gauge and gravitational theories, statistical physics and relativistic optics in locally anisotropic media and other topics) were considered in a number of works, see references from [10,11]. It seems likely that classical and quantum physical models admit a straightforward extension of la-spaces with compatible metric and connections structures. In this case we can define locally anisotropic spinors [12], formulate the theory of locally anisotropic interactions of gauge and gravitational fields [13] and consider superspaces with local anisotropy [14]. Here we remark that modern Kaluza-Klein theories can be considered as physical models on la-spaces with trivial nonlinear connection structure.

There are a number of arguments for taking into account effects of possible local anisotropy of fundamental interactions. It is well known the result that a self consistent description of radiational and dissipation processes in classical and quantum field theories

requires adding of higher derivation terms (for instance, in classical electrodynamics radiation is modeled by introducing a corresponding term proportional to the third derivation in time of coordinates). Another argument for developing quantum field models on the tangent bundle is the unclosed character of quantum electrodynamics in which the renormalized amplitudes tend to ∞ with values of momenta $p \rightarrow \infty$. This problem is avoided by introducing additional suppositions, modification of fundamental principles and by extending of the theory, which are less motivated from some physical points of view. Similar problems, but more sophisticate, arises in modeling of radiational dissipation in all variants of classical and quantum (super) gravity and (supersymmetric) quantum field theories with higher derivations. It is quite possible that to study physics of the Early Universe and propose various scenarios of Kaluza-Klein compactification from higher dimensions to the four dimensional one of the space-time it is more realistic to consider theories with generic local anisotropy caused by fluctuations of quantum high-dimensional space-time "foam". The above mentioned points to the necessity to extend the geometric background of classical and quantum field theories if a careful analysis of physical processes with non-negligible beak reaction, quantum and statistical fluctuations, turbulence, random dislocations and disclinations in continuous media.

The bulk of this paper is devoted to a review of the necessary geometric machinery for formulation of gravitational and matter field theories on la-spaces, a presenting of the essential aspects (which has not so far been published in the literature) of locally anisotropic strings and a discussion of possible generation of la-gravity from string theory.

We use R.Miron and M.Anastasei [10] conventions and basic results on the geometry of la-spaces and la-gravity.

The plan of the paper is as follows. We begin with some models of la-spaces in Sec. II ,

where we present the basic results on Finsler, Lagrange and generalized Lagrange geometry, give an overview of the geometry of nonlinear connections in vector bundles used as a general model of la-space and propose a variant of locally anisotropic Einstein-Cartan theory. In Sec. III we study the nonlinear σ -model and la-string propagation by developing the d-covariant method of la-background field. Sec. IV is devoted to problems of regularization and renormalization of the locally anisotropic σ -model and a corresponding analysis of one- and two-loop diagrams of this model. Scattering of la-gravitons and duality are considered in Sec. V., and a summary and conclusions are drawn in Sec. VI.

II. MODELS OF LOCALLY ANISOTROPIC SPACES

We present a brief introduction into the geometry of vector and tangent bundles (in brief, v-bundles and t-bundles) provided with nonlinear and distinguished connections and metric structures [10].

A. Finsler, Lagrange and Generalized Lagrange Spaces

Let M be a differentiable manifold of dimension n , $\dim = n$ (differentiable means the class C^∞ differentiability of functions) and TM is its tangent bundle. We denote by

$$u = (x, y) = \{u^\alpha = (x^i, y^{(j)} = (x^i, y^j)\},$$

$(i, j, k, \dots = n - 1)$, the local coordinates on TM where $x = (x^i)$ are local coordinates on M and $y = (y^i) = (y^{(j)})$ are fiber coordinates.

A function $\Lambda : TM \rightarrow \mathcal{R}$ ($\Lambda : (x, y) \rightarrow \Lambda(x, y)$, $\mathcal{R}$ is the real number field) is called a Finsler metric on M if it satisfies properties:

1. $\Lambda(x, y) > 0$, if $y \neq 0$;
2. Λ is a function of class C^∞ for every $y \neq 0$;
3. Λ is of homogeneity 1 on y , i.e. $\Lambda(x, \zeta y) = \zeta \Lambda(x, y)$, $\zeta \in \mathcal{R}$;
4. the quadratic form

$$g_{ij}(x, y) = \frac{1}{2} \frac{\partial^2 \Lambda^2}{\partial y^i \partial y^j}, \quad (2.1)$$

called the locally anisotropic metric tensor, la-metric, on $\mathcal{R}^n$ is positive. Couple $F^n = (M, \Lambda(x, y))$ defines a Finsler space with the fundamental function (metric) $\Lambda(x, y)$.

The length of a curve $c : \gamma \in [0, 1] \rightarrow x(\gamma) \in M$ on F^n is given by the integral

$$s = \int_0^1 \Lambda(x(\tau), \frac{dx(\tau)}{d\tau}) d\tau.$$

We point out that Riemann spaces with metrics of type $g_{ij}(x)$ are a particular class of Finsler spaces with the fundamental function of the form

$$\Lambda(x, y) = \sqrt{g_{ij}(x) y^i y^j}$$

La-metrics of type (2.1) were introduced by P. Finsler as a proposal of generalization of Riemann geometry with the aim to some applications in thermodynamics and anisotropic media physics. Variables $y = (y^j)$ can be interpreted as parameters of local anisotropy of space-time or as fluctuation parameters induced by classical and quantum field interactions in locally anisotropic media with weak radiational reaction, turbulent and stochastic processes.

The next step in extending the concept of la-space was the definition of Lagrange spaces [15] used as a geometric background for non homogeneous locally anisotropic processes [10]:

A Lagrangian on M it is a differentiable function $\mathcal{L} : TM \rightarrow \mathcal{R}$ (locally defined as $\mathcal{L} : (x, y) \rightarrow \mathcal{L}(x, y)$ and called the fundamental function) with the property that fundamental tensor, la-metric,

$$g_{ij}(x, y) = \frac{1}{2} \frac{\partial^2 \mathcal{L}}{\partial y^i \partial y^j} \quad (2.2)$$

is non degenerate.

Couple $L^n = (M, \mathcal{L})$ defines a Lagrange space.

It was also possible [10] to construct a consistent geometric theory for spaces provided with a general fundamental tensor, la-metric (not obligatory represented as a second order partial derivation of some fundamental functions as in (2.1) or (2.2)).

Couple $M^n = (M, g_{ij}(x, y))$ with a general nondegenerate tensor g_{ij} defines a generalized Lagrange space, GL-space.

In a more general form the geometry of la-spaces can be formulated in the frame of vector and tangent bundles provided with nonlinear and distinguished linear connections and metric structures

B. Nonlinear Connection in Vector Bundles

Let $\mathcal{E} = (E, \pi, F, Gr, M)$ be a locally trivial v-bundle, where $F = \mathcal{R}^m$ is the typical vector space, $\dim = m$, the structural group is taken as $Gr = GL(m, \mathcal{R})$, where $GL(m, \mathcal{R})$ is the group of linear transforms of $\mathcal{R}^m$. We locally parameterize $\mathcal{E}$ by coordinates $u^\alpha = (x^i, y^a)$, where $i, j, k, l, m, \dots = 0, 1, \dots, n-1$ and $a, b, c, d, \dots = 1, 2, \dots, m$.

Coordinate transforms $(x^k, y^a) \rightarrow (x^{k'}, y^{a'})$ on $\mathcal{E}$, considered as a differentiable manifold, are given by formulas

$$x^{k'} = x^{k'}(x^k), y^{a'} = M_a^{a'}(x)y^a, \quad (2.3)$$

where $rank(\frac{\partial x^{k'}}{\partial x^k}) = n$ and $M_a^{a'}(x) \in Gr$.

One of the fundamental objects in the geometry of la-spaces is the nonlinear connection, in brief N-connection. The first global definition of N-connection was given by W. Barthel [16] (a detailed study of N-connection structures in v-bundles and basic references are contained in [10]). Here we introduce the N-connection as a global decomposition of v-bundle $\mathcal{E}$ into horizontal, $\mathcal{HE}$, and vertical, $\mathcal{VE}$, subbundles of the tangent bundle $\mathcal{TE}$:

$$\mathcal{TE} = \mathcal{HE} \oplus \mathcal{VE}. \quad (2.4)$$

With respect to a N-connection in $\mathcal{E}$ one defines a covariant derivation operator

$$\nabla_Y A = Y^i \left\{ \frac{\partial A^a}{\partial x^i} + N_i^a(x, A) \right\} s_a, \quad (2.5)$$

where s_a are local linear independent sections of $\mathcal{E}$, $\Lambda = \Lambda^a s_a$ and $Y = Y^i s_i$ is the decomposition of vector field Y on local basis s_i on M . Differentiable functions $N_i^a(x, y)$ from (2.5) (written as functions on x^i and y^a) are called as coefficients N-connection. We have these transformation laws for components N_i^a under coordinate transforms (2.3) :

$$N_{i'}^{a'} \frac{\partial x^{i'}}{\partial x^i} = M_a^{a'} N_i^a + \frac{\partial M_a^{a'}}{\partial x^i} y^a.$$

N-connection is also characterized by its curvature

$$\Omega = \frac{1}{2} \Omega_{ij}^a dx^i \wedge dx^j \otimes \frac{\partial}{\partial y^a},$$

where $\wedge$ is the antisymmetric tensor product, with coefficients

$$\Omega_{ij}^a = \frac{\partial N_j^a}{\partial x^i} - \frac{\partial N_i^a}{\partial x^j} + N_j^b \frac{\partial N_i^a}{\partial y^b} - N_i^b \frac{\partial N_j^a}{\partial y^b},$$

and by its linearization defined as

$$\Gamma_{.bi}^a(x, y) = \frac{\partial N_i^a(x, y)}{\partial y^b}.$$

The usual linear connections

$$\omega_b^a = K_{.bi}^a(x) dx^i$$

in v-bundle $\mathcal{E}$ form a particular class of N-connections with coefficients parameterized as

$$N_i^a(x, y) = K_{.bi}^a(x) y^b.$$

C. Distinguished Tensor Fields and Connections

If in v-bundle $\mathcal{E}$ a N-connection structure is fixed, we must modify the operation of partial derivation and to introduce a locally adapted, to the N-connection, basis (frame)

$$\frac{\delta}{\delta u^\alpha} = (\frac{\delta}{\delta x^i} = \partial_i - N_i^a(x, y) \frac{\partial}{\partial y^a}, \frac{\delta}{\delta y^a} = \frac{\partial}{\partial y^a}), \quad (2.6)$$

instead of local coordinate basis

$$\frac{\partial}{\partial u^a} = (\frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^a}).$$

The basis dual to $\frac{\delta}{\delta u^\alpha}$ is written as

$$\delta u^\alpha = (\delta x^i = dx^i, \delta y^a = dy^a + N_i^a(x, y) dx^i). \quad (2.7)$$

By using bases(2.6) and (2.7) we can introduce the algebra of tensor distinguished fields (d-fields, d-tensors) on $\mathcal{E}$, $\mathcal{C} = \mathcal{C}_{qs}^{pr}$, which is equivalent to the tensor algebra of the v-bundle $\mathcal{E}_d$ defined as

$$\pi_d : \mathcal{HE} \oplus \mathcal{VE} \rightarrow \mathcal{TE}.$$

An element $t \in \mathcal{C}_{qs}^{pr}$, d-tensor of type $\begin{pmatrix} p & r \\ q & s \end{pmatrix}$, can be written in local form as

$$t = t_{j_1 \dots j_q b_1 \dots b_s}^{i_1 \dots i_p a_1 \dots a_r}(x, y) \frac{\delta}{\delta x^{i_1}} \otimes \dots \otimes \frac{\delta}{\delta x^{i_r}} \otimes dx^{j_1} \otimes \dots \otimes dx^{j_p} \otimes \frac{\partial}{\partial y^{a_1}} \otimes \dots \otimes \frac{\partial}{\partial y^{a_r}} \otimes \delta y^{b_1} \otimes \dots \otimes \delta y^{b_s}.$$

In addition to d-tensors we can consider d-objects with differential properties under group and coordinate transforms adapted to a global splitting (2.4).

A distinguished linear connection, in brief d-connection, is defined as a linear connection D in $\mathcal{E}$ conserving as a parallelism the Whitney sum $\mathcal{HE} \oplus \mathcal{VE}$ associated to a fixed N-connection structure in $\mathcal{E}$.

Components $\Gamma_{\cdot\beta\gamma}^\alpha$ of a d-connection D are introduced as

$$D_\gamma\left(\frac{\delta}{\delta u^\beta}\right) = D_{\left(\frac{\delta}{\delta u^\gamma}\right)}\left(\frac{\delta}{\delta u^\beta}\right) = \Gamma_{\cdot\beta\gamma}^\alpha\left(\frac{\delta}{\delta u^\alpha}\right). \quad (2.8)$$

We can define in a standard manner, with respect to locally adapted frame (2.6), the components of torsion $T_{\cdot\beta\gamma}^\alpha$ and curvature $R_{\beta\cdot\gamma\tau}^\alpha$ of d-connection D :

$$T\left(\frac{\delta}{\delta u^\gamma}, \frac{\delta}{\delta u^\beta}\right) = D_{\left(\frac{\delta}{\delta u^\gamma}\right)}\frac{\delta}{\delta u^\beta} - D_{\left(\frac{\delta}{\delta u^\beta}\right)}\frac{\delta}{\delta u^\gamma} - \left[\frac{\delta}{\delta u^\gamma}, \frac{\delta}{\delta u^\beta}\right] = T_{\cdot\beta\gamma}^\alpha \frac{\delta}{\delta u^\alpha},$$

where

$$T_{\cdot\beta\gamma}^\alpha = \Gamma_{\cdot\beta\gamma}^\alpha - \Gamma_{\cdot\gamma\beta}^\alpha + w_{\cdot\beta\gamma}^\alpha \quad (2.9)$$

and

$$R\left(\frac{\delta}{\delta u^\delta}, \frac{\delta}{\delta u^\gamma}, \frac{\delta}{\delta u^\beta}\right) =$$

$$(D_{(\frac{\delta}{\delta u^\delta})} D_{(\frac{\delta}{\delta u^\gamma})} - D_{(\frac{\delta}{\delta u^\gamma})} D_{(\frac{\delta}{\delta u^\delta})} - D_{([\frac{\delta}{\delta u^\delta}, \frac{\delta}{\delta u^\gamma}]}) \frac{\delta}{\delta u^\beta} = R_{\beta, \gamma \delta}^\alpha \frac{\delta}{\delta u^\alpha},$$

where

$$R_{\beta, \gamma \delta}^\alpha = \frac{\delta \Gamma_{\beta \gamma}^\alpha}{\delta u^\delta} - \frac{\delta \Gamma_{\beta \delta}^\alpha}{\delta u^\gamma} + \Gamma_{\beta \gamma}^\varphi \Gamma_{\varphi \delta}^\alpha - \Gamma_{\beta \delta}^\varphi \Gamma_{\varphi \gamma}^\alpha + \Gamma_{\beta \varphi}^\alpha w_{\gamma \delta}^\varphi. \quad (2.10)$$

In formulas (2.9) and (2.10) we have introduced nonholonomy coefficients $w_{\beta \gamma}^\alpha$ of locally adapted frames (2.6):

$$[\frac{\delta}{\delta u^\alpha}, \frac{\delta}{\delta u^\beta}] = \frac{\delta}{\delta u^\alpha} \frac{\delta}{\delta u^\beta} - \frac{\delta}{\delta u^\beta} \frac{\delta}{\delta u^\alpha} = w_{\alpha \beta}^\gamma \frac{\delta}{\delta u^\gamma}. \quad (2.11)$$

By straightforward calculations we can verify that distinguished torsion (2.9) and curvature (2.10) satisfy Bianchi identities:

$$\sum_{[\alpha \beta \gamma]} (D_\alpha R_{[\delta] \beta \gamma}^\varphi + R_{\delta, \alpha [\psi]}^\varphi T_{\beta \gamma]}^\psi) = 0,$$

$$\sum_{[\alpha \beta \gamma]} (D_\alpha T_{\beta \gamma}^\delta + T_{\alpha \beta}^\varphi T_{\gamma \varphi}^\delta - R_{\alpha, \beta \gamma}^\delta) = 0,$$

where $\sum_{[\alpha \beta \gamma]}$ denotes the antisymmetric cyclic summation on indices α, β and γ .

The global decomposition (2.4) induces a corresponding invariant splitting into horizontal $D_X^h = D_{hX}$ (h-derivation) and vertical $D_X^v = D_{vX}$ (v-derivation) parts of the operator of covariant derivation $D, D_X = D_X^h + D_X^v$, where $hX = X^i \frac{\delta}{\delta u^i}$ and $vX = X^a \frac{\partial}{\partial y^a}$ are, respectively, the horizontal and vertical components of vector field $X = hX + vX$ on $\mathcal{E}$.

Local coefficients $(L_{jk}^i(x, y), L_{bk}^a(x, y))$ of covariant h-derivation D^h are introduced as

$$D_{(\frac{\delta}{\delta x^k})}^h \left(\frac{\delta}{\delta x^j} \right) = L_{jk}^i(x, y) \frac{\delta}{\delta x^i},$$

$$D_{\left(\frac{\delta}{\delta x^k}\right)}^h \left(\frac{\partial}{\partial y^b} \right) = L_{.bk}^a(x, y) \frac{\partial}{\partial y^a}$$

and

$$D_{\left(\frac{\delta}{\delta x^k}\right)}^h f = \frac{\delta f}{\delta x^k} = \frac{\partial f}{\partial x^k} - N_k^a(x, y) \frac{\partial f}{\partial y^a}, \quad (2.12)$$

where $f(x, y)$ is a scalar function on $\mathcal{E}$.

Local coefficients $(C_{.jk}^i(x, y), C_{.bk}^a(x, y))$ of v-derivation D^v are introduced as

$$D_{\left(\frac{\partial}{\partial y^c}\right)}^v \left(\frac{\delta}{\delta x^j} \right) = C_{.jk}^i(x, y) \frac{\delta}{\delta x^i},$$

$$D_{\left(\frac{\partial}{\partial y^c}\right)}^v \left(\frac{\partial}{\partial y^b} \right) = C_{.bc}^a(x, y)$$

and

$$D_{\left(\frac{\partial}{\partial y^c}\right)}^v f = \frac{\partial f}{\partial y^c}. \quad (2.13)$$

Let us consider, for example, the action of a covariant d-derivation on a d-tensor field of type $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$

$$Q = Q_{jb}^{ia} \frac{\delta}{\delta x^i} \otimes \frac{\partial}{\partial y^a} \otimes dx^i \otimes \delta y^b.$$

We write

$$D_X Q = D_X^h Q + D_X^v Q,$$

where

$$D_X^h Q = X^k Q_{jb|k}^{ia} \frac{\delta}{\delta x^i} \otimes \frac{\partial}{\partial y^a} \otimes dx^i \otimes \delta y^b,$$

$$Q_{ib|k}^{ia} = \frac{\delta Q_{jb}^{ia}}{\delta x^k} + L_{.hk}^i Q_{jb}^{ha} - L_{.jk}^h Q_{kb}^{ia} + L_{.ck}^a Q_{jb}^{ic} - L_{.bk}^c Q_{jc}^{ia},$$

and

$$D_X^v Q = X^c Q_{jb\perp c}^{ia} \frac{\delta}{\delta x^i} \otimes \frac{\partial}{\partial y^a} \otimes dx^i \otimes \delta y^b,$$

$$Q_{jb\perp c}^{ia} = \frac{\partial Q_{jb}^{ia}}{\partial y^c} + C_{.kc}^i Q_{jb}^{ka} - C_{.jc}^k Q_{kb}^{ia} + C_{.db}^a Q_{jb}^{id} - C_{.bc}^d Q_{jd}^{ia}.$$

In order to computer the h- and v-components of torsion $T_{\beta\gamma}^\alpha$ and curvature $R_{\beta.\gamma\delta}^\alpha$ we denote :

$$hT\left(\frac{\delta}{\delta x^k}, \frac{\delta}{\delta x^j}\right) = T_{.jk}^i \frac{\delta}{\delta x^k}, hT\left(\frac{\partial}{\partial y^b}, \frac{\delta}{\delta x^i}\right) = P_{.jb}^i \frac{\delta}{\delta x^i},$$

$$vT\left(\frac{\delta}{\delta x^k}, \frac{\delta}{\delta x^j}\right) = T_{.jk}^a \frac{\partial}{\partial y^a}, vT\left(\frac{\partial}{\partial y^b}, \frac{\delta}{\delta x^i}\right) = P_{.jb}^a \frac{\partial}{\partial y^a},$$

$$vT\left(\frac{\partial}{\partial y^a}, \frac{\partial}{\partial y^b}\right) = S_{.bc}^a \frac{\partial}{\partial y^a}$$

and

$$R\left(\frac{\delta}{\delta x^k}, \frac{\delta}{\delta x^j}, \frac{\delta}{\delta x^i}\right) = R_{l.jk}^i \frac{\delta}{\delta x^i}, R\left(\frac{\delta}{\delta x^i}, \frac{\delta}{\delta x^j}, \frac{\partial}{\partial y^a}\right) = R_{b.jk}^a \frac{\partial}{\partial y^a},$$

$$R\left(\frac{\partial}{\partial y^c}, \frac{\delta}{\delta x^k}, \frac{\delta}{\delta x^j}\right) = P_{j.kc}^i \frac{\delta}{\delta x^i}, R\left(\frac{\partial}{\partial y^c}, \frac{\delta}{\delta x^k}, \frac{\partial}{\partial y^b}\right) = P_{b.kc}^a \frac{\partial}{\partial y^a},$$

$$R\left(\frac{\partial}{\partial y^c}, \frac{\partial}{\partial y^b}, \frac{\delta}{\delta x^j}\right) = S_{j.bc}^i \frac{\delta}{\delta x^i}, R\left(\frac{\partial}{\partial y^d}, \frac{\partial}{\partial y^c}, \frac{\partial}{\partial y^b}\right) = S_{b.cd}^a \frac{\partial}{\partial y^a}.$$

Putting expressions (2.12) and (2.13) into (2.9) and (2.10), after straightforward calculations, we obtain the h- and v-components of torsion,

$$T_{.jk}^i = T_{jk}^i, T_{ja}^i = C_{.ja}^i, T_{aj}^i = -C_{ja}^i, T_{.ja}^i = 0, T_{.bc}^a = S_{.bc}^a,$$

$$T_{.ij}^a = \frac{\delta N_i^a}{\delta x^j} - \frac{\delta N_j^a}{\delta x^i}, T_{.bi}^a = P_{.bi}^a = \frac{\partial N_i^a}{\partial y^b} - L_{.bj}^a, T_{.ib}^a = -P_{.bi}^a, \quad (2.14)$$

and of curvature

$$R_{h.jk}^i = \frac{\delta L_{.hj}^i}{\delta x^h} - \frac{\delta L_{.hk}^i}{\delta x^j} + L_{.hj}^m L_{mk}^i - L_{.hk}^m L_{mj}^i + C_{.ha}^i R_{.jk}^a,$$

$$\begin{aligned}
R_{b.jk}^a &= \frac{\delta L_{.bj}^a}{\delta x^k} - \frac{\delta L_{.bk}^a}{\delta x^j} + L_{.bj}^c L_{.ck}^a - L_{.bk}^c L_{.cj}^a + C_{.bc}^a R_{.jk}^c, \\
P_{j.ka}^i &= \frac{\partial L_{.jk}^i}{\partial y^k} - \left(\frac{\partial C_{.ja}^i}{\partial x^k} + L_{.lk}^i C_{.ja}^l - L_{.jk}^l C_{.la}^i - L_{.ak}^c C_{.jc}^i \right) + C_{.jb}^i P_{.ka}^b, \\
P_{b.ka}^c &= \frac{\partial L_{.bk}^c}{\partial y^a} - \left(\frac{\partial C_{.ba}^c}{\partial x^k} + L_{.dk}^c C_{.ba}^d - L_{.bk}^d C_{.da}^c - L_{.ak}^d C_{.bd}^c \right) + C_{.bd}^c P_{.ka}^d, \\
S_{j.bc}^i &= \frac{\partial C_{.jb}^i}{\partial y^c} - \frac{\partial C_{.jc}^i}{\partial y^b} + C_{.jb}^h C_{.hc}^i - C_{.jc}^h C_{.hb}^i, \\
S_{b.cd}^a &= \frac{\partial C_{.bc}^a}{\partial y^d} - \frac{\partial C_{.bd}^a}{\partial y^c} + C_{.bc}^e C_{.ed}^a - C_{.bd}^e C_{.ec}^a.
\end{aligned} \tag{2.15}$$

The components of the Ricci d-tensor

$$R_{\alpha\beta} = R_{\alpha.\beta\tau}^{\tau}$$

with respect to locally adapted frame (2.7) are as follows:

$$\begin{aligned}
R_{ij} &= R_{i.jk}^k, R_{ia} = -^2 P_{ia} = -P_{i.ka}^k, \\
R_{ai} &= ^1 P_{ai} = P_{a.ib}^b, R_{ab} = S_{a.bc}^c.
\end{aligned} \tag{2.16}$$

We point out that because, in general, $^1 P_{ai} \neq ^2 P_{ia}$ the Ricci d-tensor is non symmetric.

We shall also use the auxiliary torsionless d-connection $\tau_{.\beta\gamma}^\alpha$ (defining the corresponding auxiliary covariant d-derivation denoted as ∇) introduced as to satisfy relations

$$\Gamma_{.\beta\tau}^\alpha = \tau_{.\beta\tau}^\alpha + T_{.\beta\tau}^\alpha. \tag{2.17}$$

The Riemann and Ricci d-tensors of the d-connection $\tau_{.\beta\gamma}^\alpha$ are denoted respectively as $r_{\beta.\gamma\delta}^\alpha$ and $r_{\alpha\beta}$.

D. Distinguished Metric Structures in Vector Bundles

Now, we shall analyze the compatibility conditions of N- and d-connection and metric structures on v-bundle $\mathcal{E}$.

A metric field on $\mathcal{E}$, $G(u) = G_{\alpha\beta}(u) du^\alpha du^\beta$, is associated to a map

$$G(X, Y) : \mathcal{T}_u \mathcal{E} \times \mathcal{T}_u \mathcal{E} \rightarrow R,$$

parameterized by a non degenerate symmetric matrix

$$\begin{pmatrix} \hat{G}_{ij} & \hat{G}_{ia} \\ \hat{G}_{aj} & \hat{G}_{ab} \end{pmatrix}, \quad (2.18)$$

where

$$\hat{G}_{ij} = G\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}\right), \hat{G}_{ia} = G\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^a}\right)$$

and

$$\hat{G}_{ab} = G\left(\frac{\partial}{\partial y^a}, \frac{\partial}{\partial y^b}\right).$$

We choose a concordance between N-connection and G-metric structures by imposing conditions

$$G\left(\frac{\delta}{\delta x^i}, \frac{\partial}{\partial y^a}\right) = 0,$$

or, equivalently,

$$N_i^a(x, u) = \hat{G}_{ib}(x, y) \hat{G}^{ba}(x, y), \quad (2.19)$$

where $\hat{G}^{ba}(x, y)$ are found as components of the matrix $\hat{G}^{\alpha\beta}$ being the inverse to $\hat{G}_{\alpha\beta}$. In this case metric G on $\mathcal{E}$ is defined by two independent d-tensors, $g_{ij}(x, y)$ of type $\begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix}$ and

$h_{ab}(x, y)$ of type $\begin{pmatrix} 0 & 0 \\ 0 & 2 \end{pmatrix}$, and with respect to locally adapted basis (2.7) can be written as

$$G(u) = G_{\alpha\beta}(u) \delta u^\alpha \delta u^\beta = g_{ij}(x, y) dx^i \otimes dx^j + h_{ab}(x, y) \delta y^a \otimes \delta y^b. \quad (2.20)$$

D-connection $\Gamma_{\beta\gamma}^\alpha$ is compatible with d-metric structure $G(u)$ on $\mathcal{E}$ if one holds equalities

$$D_\alpha G_{\beta\gamma} = 0. \quad (2.21)$$

Having defined the d-metric (2.20) in $\mathcal{E}$ we can introduce the scalar curvature of d-connection

$$\overleftarrow{R} = G^{\alpha\beta} R_{\alpha\beta} = R + S, \quad (2.22)$$

where $R = g^{ij} R_{ij}$ and $S = h^{ab} S_{ab}$. The scalar curvature of the auxiliary torsionless d-connection (2.17) can be written in a similar manner, $\overleftarrow{r} = G^{\alpha\beta} r_{\alpha\beta}$.

Both d-metric (2.20) and N-connection on $\mathcal{E}$ define the so-called canonic d-connection, $C\Gamma(N)$ -connection [10], with locally adapted components

$${}^\circ\Gamma_{jk}^i = \left({}^\circ L_{jk}^i, {}^\circ L_{bi}^a, {}^\circ C_{jk}^i, {}^\circ C_{bc}^a \right)$$

expressed as

$$\begin{aligned} {}^\circ L_{jk}^i &= \frac{1}{2} g^{ip} \left(\frac{\delta g_{pj}}{\delta x^k} + \frac{\delta g_{pk}}{\delta x^j} - \frac{\delta g_{jk}}{\delta x^k} \right), \\ {}^\circ L_{bi}^a &= \frac{\partial N_i^a}{\partial y^b} + \frac{1}{2} h^{ac} \left(\frac{\delta h_{bc}}{\delta x^i} - \frac{\partial N_i^d}{\partial y^b} h_{dc} - \frac{\partial N_i^d}{\partial y^c} h_{db} \right), \\ {}^\circ C_{jb}^i &= \frac{1}{2} g^{ik} \frac{\partial g_{jk}}{\partial y^b}, \end{aligned}$$

$${}^\circ C_{.bc}^a = \frac{1}{2} h^{ad} \left(\frac{\partial h_{dc}}{\partial y^b} + \frac{\partial h_{db}}{\partial y^c} - \frac{\partial h_{bc}}{\partial y^d} \right).$$

Putting components of $CT(N)$ -connection ${}^\circ \Gamma_{.\beta\gamma}^\alpha$ instead of components of arbitrary d-connection $\Gamma_{.\beta\gamma}^\alpha$ into formulas (2.14),(2.15) and (2.16) we can calculate respectively the adapted coefficients of the canonic torsion ${}^\circ T_{.\beta\gamma}^\alpha$, curvature ${}^\circ R_{\beta.\gamma\delta}^\alpha$, and Ricci d-tensor ${}^\circ R_{\alpha\beta}$.

E. The Almost Hermitian Model of Generalized Lagrange Spaces

Let us fix a N-connection in tangent bundle (TM, τ, M) and consider a GL-space $M^n = (M, g_{ij}(x, y))$. We note that all formulas presented in the previous subsections hold good on TM . In this case the dimension of the structural fiber is equal to the dimension of base space. Both type of indices (on the base, i,j,k,..., and on the typical fiber, (i)=i,(j)=j,(k)=k,...) run from 0 to n-1. Every compatible with metric d-connection $L\Gamma(N) = (\bar{L}_{.jk}^i, \bar{C}_{jk}^i)$ on M^n can be written with respect to a locally adapted basis as

$$\begin{aligned} \bar{L}_{.jk}^i &= L_{.jk}^i + \frac{1}{2} g^{ir} (g_{rm} T_{.jk}^m - g_{jm} T_{.rk}^m + g_{km} T_{.jr}^m), \\ \bar{C}_{.jk}^i &= C_{.jk}^i + \frac{1}{2} g^{ir} (g_{im} S_{.jk}^m - g_{jm} S_{.rk}^m + g_{km} S_{.jr}^m) \end{aligned} \quad (2.23)$$

with arbitrary prescribed torsion tensors

$$T_{.jk}^i = -T_{.kj}^i, \text{ and } S_{.ij}^i = -S_{.kj}^i.$$

Coefficients

$$\begin{aligned} L_{.jk}^i(x, y) &= \frac{1}{2} g^{ir} \left(\frac{\delta g_{jk}}{\delta x^k} + \frac{\delta g_{kr}}{\delta x^j} - \frac{\delta g_{jk}}{\delta x^r} \right), \\ C_{.jk}^i(x, y) &= \frac{1}{2} g^{ir} \left(\frac{\partial g_{jr}}{\partial y^k} + \frac{\partial g_{kr}}{\partial y^j} - \frac{\partial g_{jk}}{\partial y^r} \right) \end{aligned} \quad (2.24)$$

form the unique torsionless canonic connection $C\Gamma(N) = (L^i_{.jk}, C^i_{.jk})$ on M^n .

N-connection defines naturally an almost complex structure [10] F on TM acting as

$$F\left(\frac{\delta}{\delta x^i}\right) = -\frac{\partial}{\partial y^i},$$

and

$$F\left(\frac{\partial}{\partial y^i}\right) = \frac{\delta}{\delta x^i}.$$

If an almost tangent structure J , with the property that

$$J\left(\frac{\delta}{\delta x^i}\right) = \frac{\delta}{\delta x^i}$$

and

$$J\left(\frac{\partial}{\partial y^i}\right) = -\frac{\partial}{\partial y^i},$$

and a d-connection D satisfying condition $DJ = 0$ are defined on TM , one calls D the normal connection on M^n .

One says that a linear connection D on TM is the N- lift of a normal d-connection on M if the compatibility of the almost complex structure F and of d-connection structure holds, i.e. if $DF = 0$.

Condition $g(FX, FY) = g(X, Y)$, where X and Y are arbitrary d-vector fields, for d-metric (2.20) on TM implies that $g_{ij}(x, y) = g_{(i)(j)}(x, y)$, i.e. the N-lift on TM of the fundamental tensor $g_{ij}(x, y)$ on M defines a covariant second order symmetric non degenerate d-field

$$\vec{G}(x, y) = g_{ij}(x, y) dx^i dx^j + g_{ij}(x, y) \delta y^i \delta y^j. \quad (2.25)$$

Let call this d-metric as the H^{2n} -metric. By straightforward calculations we can verify that H^{2n} -metric is compatible with the N-lift of metric d-connection $L\Gamma(N)$ (2.23), i.e.

$$\overline{D}_\gamma \overset{\leftrightarrow}{G}_{\alpha\beta} = 0. \quad (2.26)$$

Components of connection $\overline{D}$ on TM with respect to a locally adapted basis are computed [10] as

$$\begin{aligned} \overline{\Gamma}^i_{jk} &= \overline{L}^i_{jk}, \overline{\Gamma}^i_{j(k)} = \overline{C}^i_{jk}, \\ \overline{\Gamma}^{(i)}_{(j)k} &= \overline{L}^i_{jk}, \overline{\Gamma}^{(i)}_{(j)(k)} = \overline{C}^i_{jk} \end{aligned} \quad (2.27)$$

Space $H^{2n} = (TM, G, F)$, in brief H^{2n} -space, defines the almost Hermitian model of GL-space $M^n = (M, g_{ij}(x, y))$.

Components of torsion $\overline{T}^\alpha_{\beta\gamma}$, curvature and Ricci tensor $\overline{R}^\alpha_{\beta\gamma\delta}$ and Ricci tensor $\overline{R}_{\alpha\beta}$ can be computed respectively by putting components (2.27) into formulas (2.14),(2.15) and (2.16) considered on tangent bundle TM .

Finally, in this subsection, we remark that we can consider N-lifts on TM of a Finsler (or Lagrange) la-metric, if instead of coefficients of GL-metric g_{ij} the coefficients (2.1) (on (2.2)) are used.

F. Einstein-Cartan Equations on Locally Anisotropic Spaces

Now we shall analyze the problem of formulation of models of gravitational interactions on la-spaces and propose a version of la-gravity theory with torsion on such spaces.

There are developed some approaches to la-gravity (see details and discussions of physical aspects in Refs [11]). Here we emphasize that the Einstein like theory on H^{2n} - spaces [10] extends the general relativity theory on la-spaces in a manner as to have compatible nonlinear and distinguished linear connections and metric structures (see (2.26)). This property allows a straightforward generalization of classical and quantum field theories on

(pseudo)- Riemannian spaces to similar ones on H^{2n} -spaces by changing respectively partial derivations $\frac{\partial}{\partial x^i}$, metric $g_{ij}(x)$ and metric connection $\Gamma_{jk}^i(x)$ into locally adapted operators $\frac{\delta}{\delta u^\alpha} = \left(\frac{\delta}{\delta x^i}, \frac{\partial}{\partial y^j}\right)$ (see (2.6)), H^{2n} -metric $\overleftrightarrow{G}_{\mu\nu}(u)$ (see (2.25)) and H^{2n} -connection $\overline{\Gamma}^\alpha_{\beta\gamma}$ (see (2.27)). We also note that on generic H^{2n} -spaces the locally adapted to N-connection bases are nonholonomic, torsion is not vanishing and the Ricci d-tensor is not symmetric.

As a general model of la-space we shall use a v-bundle space $\mathcal{E}$ provided with N-connection, d-connection and d-metric satisfying compatibility conditions (2.21).

Einstein-Cartan equations for d-connection $\Gamma^\alpha_{\beta\gamma}$ are written in a standard manner by using the torsion d-tensor (2.6), the Ricci d-tensor (2.16), d-metric (2.20) and the scalar curvature (2.22):

$$R_{\alpha\beta} - \frac{1}{2}G_{\alpha\beta}\overleftarrow{R} + \tilde{\lambda}G_{\alpha\beta} = \kappa_{(1)}T_{\alpha\beta}, \quad (2.28)$$

$$T^\alpha_{\beta\gamma} + \delta^\alpha_\beta T^\tau_{\gamma\tau} - \delta^\alpha_\gamma T^\tau_{\beta\tau} = \kappa_{(2)}Q^\alpha_{\beta\gamma}, \quad (2.29)$$

where $\kappa_{(1)}$ and $\kappa_{(2)}$ are interaction constants. In (2.28) we have introduced the locally anisotropic cosmological term $\tilde{\lambda}G_{\alpha\beta}$ and energy-momentum d-tensor $T_{\alpha\beta}$. D-tensor $Q^\alpha_{\beta\gamma}$ from (2.29) is interpreted as the locally anisotropic spin-matter source of torsion.

R. Miron and M. Anastasiei [17,10] considered on v-bundle ξ the Einstein equations with prescribed torsion and N-connection and without cosmological term (we present here the h- and v-projections of their equations, see, correspondingly, the necessary h- and v-components of d-metric, Ricci d-tensor and scalar curvature in (2.20),(2.16) and (2.22)):

$$R_{ij} - \frac{1}{2}(R + S)g_{ij} = \kappa_{(1)}T_{ij},$$

$$S_{ab} - \frac{1}{2}(R + S)h_{ab} = \kappa_{(2)}T_{ab},$$

$${}^1P_{ai} = \kappa_{(1)}T_{ai},$$

$${}^2P_{ia} = -\kappa_{(2)}T_{ia}. \quad (2.30)$$

La-gravitational field equations (2.30) contain $(n + m)^2$ equations for

$$nm + \frac{1}{2}[n(n + 1) + m(m + 1) + n^2(n - 1) + m^2(m - 1)]$$

unknown variables $N_i^a, g_{ij}, h_{ab}, T_{jk}^i$ and S_{bc}^a . The last number is greater than the first one for $n \geq 3$ and $m \geq 1$. We have add equations (2.29) in order to obtain a closed system of equations with metric-N-connection constraints of type (2.18). The introduction of the cosmological term presents interest for a further study of possible cosmological consequences of la-gravity. In [13] we formulated a more general variant of gauge la-gravity with dynamical equations for torsion instead of algebraic equations. In this work we shall consider both the Einstein like field equations and the equations for the curvature of N-connection and torsion of d-connection arising as constraints for a consistent propagation of strings on la-backgrounds.

Finally, in this section, we note that la-gravitational field equations (2.30) will be transformed into Einstein ones for multidimensional gravity if the nonlinear connection and torsion of linear connection vanish. Usually one considers different scenarios of compactification from higher dimensions to four dimensional one by introducing by hand or postulating a kind of spontaneous symmetry breaking of local multidimensional gravitational symmetry. By using the N-connection we can modelate dynamical variants of "splitting" of space-time dimensions and consider physical consequences for possible relic effects of locally anisotropic fluctuations.

III. LOCALLY ANISOTROPIC STRINGS AND SIGMA-MODELS

In this section we present a generalization of some necessary results on nonlinear σ -model and string propagation to the case of la-backgrounds. Calculations on both type of locally anisotropic and isotropic spaces are rather similar if we accept the Miron and Anastasiei [10] geometric formalism. We emphasize that on la-backgrounds we have to take into account the distinguished character, by N-connection, of geometric objects.

A. Action of Nonlinear σ -model and Torsion of La-space

Let a map of a two-dimensional (2d), for simplicity, flat space M^2 into la-space ξ defines a σ -model field $u^\mu(z) = (x^i(z), y^a(z))$, where $z = \{z^A, A = 0, 1\}$ are two-dimensional complex coordinates on M^2 . The moving of a bosonic string in la-space is governed by the nonlinear σ -model action (see, for instance, [1,13,4] for details on locally isotropic spaces):

$$I = \frac{1}{\lambda^2} \int d^2z \left[\frac{1}{2} \sqrt{\gamma} \gamma^{AB} \partial_A u^\mu(z) \partial_B u^\nu(z) G_{\mu\nu}(u) + \frac{\tilde{n}}{3} \epsilon^{AB} \partial_A u^\mu \partial_B u^\nu b_{\mu\nu}(u) + \frac{\lambda^2}{4\pi} \sqrt{\gamma} R^{(2)} \Phi(u) \right], \quad (3.1)$$

where λ^2 and $\tilde{n}$ are interaction constants, $\Phi(u)$ is the dilaton field, $R^{(2)}$ is the curvature of the 2d world sheet provided with metric γ_{AB} , $\gamma = \det(\gamma_{AB})$ and $\partial_A = \frac{\partial}{\partial z^A}$, tensor ϵ^{AB} and d-tensor $b_{\mu\nu}$ are antisymmetric.

From the viewpoint of string theory we can interpret $b_{\alpha\beta}$ as the vacuum expectation of the antisymmetric, in our case locally anisotropic, d-tensor gauge field $B_{\alpha\beta\gamma}$ (see considerations for locally isotropic models in [18,19] and the Wess-Zumino-Witten model [20,21], which lead to the conclusion [22] that $\tilde{n}$ takes only integer values and that in the perturbative quantum field theory the effective quantum action depends only on B... and does not depend on b...).

In order to obtain compatible with N-connection motions of la-strings we consider these relations between d-tensor $b_{\alpha\beta}$, strength $B_{\alpha\beta\gamma} = \delta_{[\alpha} b_{\beta\gamma]}$ and torsion $T_{\beta\gamma}^{\alpha}$:

$$\delta_{\alpha} b_{\beta\gamma} = T_{\alpha\beta\gamma}, \quad (3.2)$$

with the integrability conditions

$$\Omega_{ij}^a \partial_a b_{\beta\gamma} = \delta_{[i} T_{j]\beta\gamma}, \quad (3.3)$$

where Ω_{ij}^a are the coefficients of the N-connection curvature. In this case we can express $B_{\alpha\beta\gamma} = T_{[\alpha\beta\gamma]}$. Conditions (3.2) and (3.3) define a simplified model of la-strings when the σ -model antisymmetric strength is induced from the la-background torsion. More general constructions are possible by using normal coordinates adapted to both N-connection and torsion structures on la-background space. For simplicity, we omit such considerations in this work.

Choosing the complex (conformal) coordinates $z = \iota^0 + i\iota^1, \bar{z} = \iota^0 - i\iota^1$, where $\iota^A, A = 0, 1$ are real coordinates, on the world sheet we can represent the two-dimensional metric in the conformally flat form:

$$ds^2 = e^{2\varphi} dz d\bar{z}, \quad (3.4)$$

where $\gamma_{z\bar{z}} = \frac{1}{2}e^{2\varphi}$ and $\gamma_{zz} = \gamma_{\bar{z}\bar{z}} = 0$.

Let us consider an la-field $U(u), u \in \xi$ taking values in $\mathcal{G}$ being the Lie algebra of a compact and semi simple Lie group,

$$U(u) = \exp[i\varphi(u)], \varphi(u) = \varphi^{\underline{\alpha}}(u) q^{\underline{\alpha}},$$

where $q^{\underline{\alpha}}$ are generators of the Lie algebra with antisymmetric structural constants $f^{\underline{\alpha}\underline{\beta}\underline{\gamma}}$ satisfying conditions

$$[q^{\underline{\alpha}}, q^{\underline{\beta}}] = 2if^{\underline{\alpha}\underline{\beta}\underline{\gamma}} q^{\underline{\gamma}}, \quad tr(q^{\underline{\alpha}} q^{\underline{\beta}}) = 2\delta^{\underline{\alpha}\underline{\beta}}.$$

The action of the Wess-Zumino-Witten type la-model should be written as

$$I(U) = \frac{1}{4\lambda^2} \int d^2z \operatorname{tr} \left(\partial_A U \partial^A U^{-1} \right) + \tilde{n} \Gamma[U], \quad (3.5)$$

where $\Gamma[U]$ is the standard topologically invariant functional [22]. For perturbative calculations in the framework of the model (3.1) it is enough to know that as a matter of principle we can represent the action of our theory as (3.5) and to use d-curvature $r_{\beta.\gamma\delta}^\alpha$ for the torsionless d-connection $\tau_{\beta\gamma}^\alpha$, see (2.17), and strength $B_{\alpha\beta\gamma}$ respectively expressed as

$$R_{\alpha\beta\gamma\delta} = f_{\underline{\alpha}\underline{\beta}\underline{\tau}} f_{\underline{\gamma}\underline{\delta}\underline{\tau}} V_{\underline{\alpha}}^\alpha V_{\underline{\beta}}^\beta V_{\underline{\gamma}}^\gamma V_{\underline{\delta}}^\delta$$

and

$$B_{\alpha\beta\gamma} = \eta f_{\underline{\alpha}\underline{\beta}\underline{\tau}} V_{\underline{\alpha}}^\alpha V_{\underline{\beta}}^\beta V_{\underline{\tau}}^\tau,$$

where a new interaction constant $\eta \equiv \frac{\tilde{n}\lambda^2}{2\pi}$ is used and $V_{\underline{\alpha}}^\alpha$ is a locally adapted vielbein associated to the metric (2.20):

$$G_{\alpha\beta} = V_{\underline{\alpha}}^\alpha V_{\underline{\beta}}^\beta \delta^{\underline{\alpha}\underline{\beta}}$$

and

$$G^{\alpha\beta} V_{\underline{\alpha}}^\alpha V_{\underline{\beta}}^\beta = \delta^{\underline{\alpha}\underline{\beta}}. \quad (3.6)$$

For simplicity, we shall omit underlining of indices if this will not give rise to ambiguities.

Finally, in this subsection, we remark that for $\eta = 1$ we obtain a conformally invariant two-dimensional quantum field theory (being similar to those developed in [23,20]).

B. The D-covariant Method of La-background Field and σ -models

Suggesting the compensation of all anomalies we can fix the gauge for the two-dimensional metric when action (3.1) is written as

$$I[u] = \frac{1}{2\lambda^2} \int d^2z \{G_{\alpha\beta}\eta^{AB} + \frac{2}{3}b_{\alpha\beta}\epsilon^{AB}\} \partial_A u^\alpha \partial_B u^\beta, \quad (3.1a)$$

where η^{AB} and ϵ^{AB} are, respectively, constant two-dimensional metric and antisymmetric tensor. The covariant method of background field, as general references see [24-27], can be extended for la-spaces. Let consider a curve in ξ parameterized as $\rho^\alpha(z, s), s \in [0, 1]$, satisfying autoparallel equations

$$\begin{aligned} \frac{d^2 \rho^\alpha(z, s)}{ds^2} + \Gamma_{\beta\gamma}^\alpha[\rho] \frac{d\rho^\beta}{ds} \frac{d\rho^\gamma}{ds} = \\ \frac{d^2 \rho^\alpha(z, s)}{ds^2} + \tau_{\beta\gamma}^\alpha[\rho] \frac{d\rho^\beta}{ds} \frac{d\rho^\gamma}{ds} = 0, \end{aligned}$$

with boundary conditions

$$\rho(z, s=0) = u(z) \text{ and } \rho(z, s=1) = u(z) + v(z).$$

For simplicity, hereafter we shall consider that d-connection $\Gamma_{\beta\gamma}^\alpha$ is defined by d-metric (2.20) and N-connection structures, i.e.

$$\Gamma_{\beta\gamma}^\alpha = {}^\circ\Gamma_{\beta\gamma}^\alpha.$$

The tangent d-vector $\zeta^\alpha = \frac{d}{ds}\rho^\alpha$, where $\zeta^\alpha|_{s=0} = \zeta_{(0)}^\alpha$ is chosen as the quantum d-field. Then the expansion of action $I[u + v(\zeta)]$, see (3.1a), as power series on ζ ,

$$I[u + v(\zeta)] = \sum_{k=0}^{\infty} I_k,$$

where

$$I_k = \frac{1}{k!} \frac{d^k}{ds^k} I[\rho(s)]|_{s=0},$$

defines d-covariant, depending on the background d-field, interaction vortexes of locally anisotropic σ -model.

In order to compute I_k it is useful to consider relations

$$\frac{d}{ds}\partial_A\rho^\alpha = \partial_A\zeta^\alpha, \frac{d}{ds}G_{\alpha\beta} = \zeta^\tau\delta_\tau G_{\alpha\beta},$$

$$\partial_A G_{\alpha\beta} = \partial_A \rho^\tau \delta_\tau G_{\alpha\beta},$$

to introduce auxiliary operators

$$\begin{aligned} (\widehat{\nabla}_A \zeta)^\alpha &= (\nabla_A \zeta)^\alpha - G^{\alpha\tau} T_{[\alpha\beta\gamma]} [\rho] \epsilon_{AB} \partial^B \rho^\gamma \zeta^\beta, \\ (\nabla_A \zeta)^\alpha &= [\delta_\beta^\alpha \partial_A + \tau_{\beta\gamma}^\alpha \partial_A \rho^\gamma] \zeta^\beta, \\ \nabla(s) \xi^\lambda &= \zeta^\alpha \nabla_\alpha \xi^\lambda = \frac{d}{ds} \xi^\lambda + \tau_{\beta\gamma}^\lambda [\rho(s)] \zeta^\beta \xi^\gamma, \end{aligned} \tag{3.7}$$

having properties

$$\nabla(s) \zeta^\alpha = 0, \nabla(s) \partial_A \rho^\alpha = (\nabla_A \zeta)^\alpha,$$

$$\nabla^2(s) \partial_A \rho^\alpha = r_{\beta\gamma\delta}^\alpha \zeta^\beta \zeta^\gamma \partial_A \rho^\delta,$$

and to use the curvature d-tensor of d-connection (3.7),

$$\widehat{r}_{\beta\alpha\gamma\delta} = r_{\beta\alpha\gamma\delta} - \nabla_\gamma T_{[\alpha\beta\delta]} + \nabla_\delta T_{[\alpha\beta\gamma]} - T_{[\tau\alpha\gamma]} G^{\tau\lambda} T_{[\lambda\delta\beta]} + T_{[\tau\alpha\delta]} G^{\tau\lambda} T_{[\lambda\gamma\beta]}.$$

Values I_k can be computed in a similar manner as in [19,22,27], but in our case by using corresponding d-connections and d-objects. Here we present the first four terms in explicit form:

$$\begin{aligned} I_1 &= \frac{1}{2\lambda^2} \int d^2 z 2G_{\alpha\beta} (\widehat{\nabla}_A \zeta)^\alpha \partial^A u^\beta, \\ I_2 &= \frac{1}{2\lambda^2} \int d^2 z \{ (\widehat{\nabla}_A \zeta)^2 + \widehat{r}_{\beta\alpha\gamma\delta} \zeta^\beta \zeta^\gamma (\eta^{AB} - \epsilon^{AB}) \partial_A u^\alpha \partial_B u^\beta \}, \\ I_3 &= \frac{1}{2\lambda^2} \int d^2 z \{ \frac{4}{3} (r_{\beta\alpha\gamma\delta} - G^{\tau\epsilon} T_{[\epsilon\alpha\beta]} T_{[\tau\gamma\delta]}) \partial_A u^\alpha (\widehat{\nabla}^A \zeta^\delta) \zeta^\beta \zeta^\gamma + \frac{4}{3} \nabla_\alpha T_{[\delta\beta\gamma]} \partial_A u^\beta \epsilon^{AB} (\widehat{\nabla}_B \zeta^\gamma) \zeta^\alpha \zeta^\delta + \\ &\quad \frac{2}{3} T_{[\alpha\beta\gamma]} (\widehat{\nabla}_A \zeta^\alpha) \epsilon^{AB} (\widehat{\nabla} \zeta^\beta) \zeta^\gamma + \frac{1}{3} (\nabla_\lambda r_{\beta\alpha\gamma\delta} + 4G^{\tau\epsilon} T_{[\epsilon\lambda\alpha]} \nabla_\beta T_{[\gamma\delta\tau]}) \partial_A u^\alpha \partial^A u^\delta \zeta^\gamma \zeta^\lambda + \end{aligned}$$

$$\begin{aligned}
& \frac{1}{3} \left(\nabla_\alpha \nabla_\beta T_{[\tau\gamma\delta]} + 2G^{\lambda\epsilon} G^{\varphi\phi} T_{[\alpha\lambda\varphi]} T_{[\epsilon\beta\delta]} T_{[\phi\tau\gamma]} + 2r_{\alpha,\beta\gamma}^\lambda T_{[\alpha\tau\delta]} \right) \partial_A u^\gamma \epsilon^{AB} (\partial_B \zeta^\tau) \zeta^\alpha \zeta^\beta \}, \\
& I_4 = \frac{1}{4\lambda^2} \int d^2 z \{ \left(\frac{1}{2} \nabla_\alpha r_{\gamma\beta\delta\tau} - G^{\lambda\epsilon} T_{[\epsilon\beta\gamma]} \nabla_\alpha T_{[\lambda\delta\tau]} \right) \partial_A u^\beta (\widehat{\nabla}^A \zeta^\tau) \zeta^\alpha \zeta^\gamma \zeta^\delta + \\
& \frac{1}{3} r_{\beta\alpha\gamma\delta} (\widehat{\nabla}_A \zeta^\alpha) (\widehat{\nabla}^A \zeta^\delta) \zeta^\beta \zeta^\gamma + \left(\frac{1}{12} \nabla_\alpha \nabla_\beta r_{\delta\gamma\tau\lambda} + \frac{1}{3} r_{\delta,\tau\gamma}^\kappa r_{\beta\kappa\alpha\lambda} - \frac{1}{2} (\nabla_\alpha \nabla_\beta T_{[\gamma\tau\epsilon]}) G^{\epsilon\pi} T_{[\pi\delta\lambda]} - \right. \\
& \left. \frac{1}{2} r_{\alpha,\beta\gamma}^\kappa T_{[\kappa\tau\epsilon]} G^{\epsilon\pi} T_{[\pi\delta\lambda]} + \frac{1}{6} r_{\alpha,\beta\epsilon}^\kappa T_{[\kappa\gamma\tau]} G^{\epsilon\pi} T_{[\epsilon\delta\lambda]} \right) \partial_A u^\gamma \partial^A u^\lambda \zeta^\alpha \zeta^\beta \zeta^\delta \zeta^\tau + \left[\frac{1}{12} \nabla_\alpha \nabla_\beta \nabla_\gamma T_{[\lambda\delta\tau]} + \right. \\
& \left. \frac{1}{2} \nabla_\alpha \left(G^{\kappa\epsilon} T_{[\epsilon\lambda\delta]} \right) r_{\beta\kappa\gamma\tau} + \frac{1}{2} \left(\nabla_\alpha T_{[\pi\beta\kappa]} \right) G^{\pi\epsilon} G^{\kappa\nu} T_{[\epsilon\gamma\delta]} T_{[\nu\lambda\tau]} - \frac{1}{3} G^{\kappa\epsilon} T_{[\epsilon\lambda\delta]} \nabla_\alpha r_{\beta\kappa\delta\tau} \right] \times \\
& \partial_A u^\delta \epsilon^{AB} \partial_B u^\tau \zeta^\lambda \zeta^\alpha \zeta^\beta \zeta^\gamma + \frac{1}{2} \left[\nabla_\alpha \nabla_\beta T_{[\tau\gamma\delta]} + T_{[\kappa\delta\tau]} r_{\alpha,\beta\gamma}^\kappa + r_{\alpha,\beta\delta}^\kappa T_{[\kappa\gamma\tau]} \right] \times \\
& \partial_A u^\gamma \epsilon^{AB} (\widehat{\nabla}_B \zeta^\delta) \zeta^\alpha \zeta^\beta \zeta^\tau + \frac{1}{2} \nabla_\alpha T_{[\delta\beta\gamma]} (\widehat{\nabla}_A \zeta^\beta) \epsilon^{AB} (\widehat{\nabla}_B \zeta^\gamma) \zeta^\alpha \zeta^\delta \}. \tag{3.8}
\end{aligned}$$

Now we construct the d-covariant la-background functional (we use methods, in our case correspondingly adapted to the N-connection structure, developed in [19,27,28]). The standard quantization technique is based on the functional integral

$$Z[J] = \exp(iW[J]) = \int d[u] \exp\{i(I + uJ)\}, \tag{3.9}$$

with source J^α (we use condensed denotations and consider that computations are made in the Euclidean space). The generation functional Γ of one-particle irreducible (1PI) Green functions is defined as

$$\Gamma[\bar{u}] = W[J(\bar{u})] - \bar{u} \cdot J[\bar{u}],$$

where $\bar{u} = \frac{\Delta W}{\delta J}$ is the mean field. For explicit perturbative calculations it is useful to connect the source only with the covariant quantum d-field ζ and to use instead of (3.9) the new functional

$$\exp(iW[\bar{u}, J]) = \int [d\zeta] \exp\{i(I[\bar{u} + v(\zeta)] + J \cdot \zeta)\}. \tag{3.10}$$

It is clear that Feynman diagrams obtained from this functional are d-covariant.

Defining the mean d-field $\bar{\zeta}(u) = \frac{\Delta W}{\delta J(u)}$ and introducing the auxiliary d-field $\zeta' = \zeta - \bar{\zeta}$ we obtain from (3.9) a double expansion on both classical and quantum la-backgrounds:

$$\exp(i\bar{\Gamma}[\bar{u}, \bar{\zeta}]) = \int [d\zeta'] \exp\{i \left(I [\bar{u} + v(\zeta' + \bar{\zeta})] - \zeta' \frac{\Delta \Gamma}{\delta \bar{\zeta}} \right)\}. \quad (3.11)$$

The manner of fixing the measure in the functional (3.10) (and as a consequence in (3.11)) is obvious :

$$[d\zeta] = \prod_u \sqrt{|G(u)|} \prod_{\alpha=0}^{n+m-1} d\zeta^\alpha(u). \quad (3.12)$$

Using vielbein fields (3.6) we can rewrite the measure (3.12) in the form

$$[d\zeta] = \prod_u \prod_{\alpha=0}^{n+m-1} d\zeta^\alpha(u).$$

The structure of renormalization of σ -models of type (3.10) (or (3.11)) is analyzed, for instance, in [19,27,28]. For la-spaces we must take into account the N-connection structure.

IV. REGULARIZATION AND β -FUNCTIONS OF THE LOCALLY ANISOTROPIC σ -MODEL

The aim of this section is to study the problem of regularization and quantum ambiguities in β -functions of the renormalization group and to present the results on one- and two-loop calculus for the la- σ -model (LAS-model).

A. Regularization and Renormalization Group β -functions

Because our σ -model is a two-dimensional and massless locally anisotropic theory we have to consider both types of infrared and ultraviolet regularizations (in brief, IR- and UV-regularization). In order to regularize IR-divergences and distinguish them from UV-divergences we can use a standard mass term in the action (3.1) of the LAS-model

$$I_{(m)} = -\frac{\tilde{m}^2}{2\lambda^2} \int d^2z G_{\alpha\beta} u^\alpha u^\beta.$$

For regularization of UV-divergences it is convenient to use the dimensional regularization. For instance, the regularized propagator of quantum d-fields ζ looks like

$$\begin{aligned} \langle \zeta^\alpha(u_1) \zeta^\beta(u_2) \rangle &= \delta^{\alpha\beta} G(u_1 - u_2) = \\ &= i\lambda^2 \delta^{\alpha\beta} \int \frac{d^q p}{(2\pi)^q} \frac{\exp[-ip(u_1 - u_2)]}{(p^2 - \tilde{m}^2 + i0)}, \end{aligned}$$

where $q = 2 - 2\epsilon$.

The d-covariant dimensional regularization of UV-divergences is complicated because of existence of the antisymmetric symbol ϵ^{AB} . One introduces [29,27] this general prescription:

$$\epsilon^{LN} \eta_{NM} \epsilon^{MR} = \psi(\epsilon) \eta^{LR}$$

and

$$\epsilon^{MN} \epsilon^{RS} = \omega(\epsilon) [\eta^{MS} \eta^{NR} - \eta^{MR} \eta^{NS}],$$

where η^{MN} is the q-dimensional Minkowski metric, and $\psi(\epsilon)$ and $\omega(\epsilon)$ are arbitrary d-functions satisfying conditions $\psi(0) = \omega(0) = 1$ and depending on the type of renormalization.

We use the standard dimensional regularization, with dimensionless scalar d-field $u^\alpha(z)$, when expressions for unrenormalized $G_{\alpha\beta}^{(ur)}$ and $B_{\alpha\beta}^{(k,l)}$ have a d-tensor character, i.e. they

are polynoms on d-tensors of curvature and torsion and theirs d-covariant derivations (for simplicity in this subsection we consider $\lambda^2 = 1$; in general one-loop 1PI-diagrams must be proportional to $(\lambda^2)^{l-1}$).

RG β -functions are defined by relations (for simplicity we shall omit index R for renormalized values)

$$\begin{aligned}\mu \frac{d}{d\mu} G_{\alpha\beta} &= \beta_{(\alpha\beta)}^G (G, B), \mu \frac{d}{d\mu} B_{[\alpha\beta]} = \beta_{[\alpha\beta]}^B (G, B), \\ \beta_{\alpha\beta} &= \beta_{(\alpha\beta)}^G + \beta_{[\alpha\beta]}^B.\end{aligned}$$

By using the scaling property of the one-loop counter-term under global conformal transforms

$$G_{\alpha\beta}^G \rightarrow \Lambda^{(l-1)} G_{\alpha\beta}^{(k,l)}, B_{\alpha\beta}^{(k,l)} \rightarrow \Lambda^{(l-1)} B_{\alpha\beta}^{(k,l)}$$

we obtain

$$\beta_{(\alpha\beta)}^G = - \sum_{l=1}^{(1,l)} l G_{(\alpha\beta)}^{(1,l)}, \beta_{[\alpha\beta]}^B = - \sum_{l=1}^{\infty} l B_{[\alpha\beta]}^{(1,l)}$$

in the leading order on ϵ (compare with the usual perturbative calculus from [31]).

The d-covariant one-loop counter-term is taken as

$$\Delta I^{(l)} = \frac{1}{2} \int d^2 z T_{\alpha\beta}^{(l)} (\eta^{AB} - \epsilon^{AB}) \partial_A u^\alpha \partial_B u^\beta,$$

where

$$T_{\alpha\beta}^{(l)} = \sum_{k=1}^l \frac{1}{(2\epsilon^k)} T_{\alpha\beta}^{(k,l)} (G, B). \quad (4.1)$$

For instance, in the three-loop approximation we have

$$\beta_{\alpha\beta} = T_{\alpha\beta}^{(1,1)} + 2T_{\alpha\beta}^{(1,2)} + 3T_{\alpha\beta}^{(1,3)}. \quad (4.2)$$

In the next subsection we shall also consider constraints on the structure of β -functions connected with conditions of integrability (caused by conformal invariance of the two-dimensional world-sheet).

B. One-loop Divergences and RG-equations of the LAS-model

We generalize the one-loop results [32] to the case of la-backgrounds. If in locally isotropic models one considers an one-loop diagram, for the LAS-model the distinguished by N-connection character of la-interactions leads to the necessity to consider four one-loop diagrams (see Fig. 1). To these diagrams one corresponds counter-terms:

$$\begin{aligned}
I_1^{(c)} &= I_1^{(c,x^2)} + I_1^{(c,y^2)} + I_1^{(c,xy)} + I_1^{(c,yx)} = \\
&-\frac{1}{2}I_1 \int d^2z \hat{r}_{ij} (\eta^{AB} - \epsilon^{AB}) \partial_A x^i \partial_B x^j - \\
&\frac{1}{2}I_1 \int d^2z \hat{r}_{ab} (\eta^{AB} - \epsilon^{AB}) \partial_A y^a \partial_B y^b - \\
&\frac{1}{2}I_1 \int d^2z \hat{r}_{ia} (\eta^{AB} - \epsilon^{AB}) \partial_A x^i \partial_B y^a - \\
&\frac{1}{2}I_1 \int d^2z \hat{r}_{ai} (\eta^{AB} - \epsilon^{AB}) \partial_A y^a \partial_B x^i,
\end{aligned}$$

where I_1 is the standard integral

$$\begin{aligned}
I_1 &= \frac{G(0)}{\lambda^2} = i \int \frac{d^d p}{(2\pi)^2} \frac{1}{p^2 - \tilde{m}^2} = \frac{\Gamma(\epsilon)}{4\pi^{\frac{d}{2}}(\tilde{m}^2)^\epsilon} = \\
&\frac{1}{4\pi\epsilon} - \frac{1}{\pi} \ln \tilde{m} + \text{finite counter terms.}
\end{aligned}$$

There are one-loops on the base and fiber spaces or describing quantum interactions between fiber and base components of d-fields. If the la-background d-connection is of distinguished Levi-Civita type we obtain only two one-loop diagrams (on the base and in the fiber) because in this case the Ricci d-tensor is symmetric. It is clear that this four-multiplying (doubling for the Levi-Civita d-connection) of the number of one-loop diagrams is caused by the "indirect" interactions with the N-connection field. Hereafter, for simplicity, we shall use a compactified (non-distinguished on x- and y-components) form of

writing out diagrams and corresponding formulas and emphasize that really all expressions containing components of d-torsion generate four irreducible, see (2.14), types of diagrams (with respective interaction constants) and that all expressions containing components of d-curvature give rise in a similar manner to six irreducible, see formulas (2.17), types of diagrams. We shall take into consideration these details in the subsection where we shall write the two-loop effective action.

Subtracting in a trivial manner I_1 ,

$$I_1 + \text{subtractions} = \frac{1}{4\pi\epsilon},$$

we can write the one-loop β -function in the form:

$$\begin{aligned} \beta_{\alpha\beta}^{(1)} &= \frac{1}{2\pi} \hat{r}_{\alpha\beta} \\ &= \frac{1}{2\pi} \left(r_{\alpha\beta} - G_{\alpha\tau} T^{[\tau\gamma\phi]} T_{[\beta\gamma\phi]} + G^{\tau\mu} \nabla_\mu T_{[\alpha\beta\tau]} \right). \end{aligned}$$

We also note that the mass term in the action generates the mass one-loop counter-term

$$\Delta I_1^{(m)} = \frac{\tilde{m}^2}{2} I_1 \int d^2 z \left\{ \frac{1}{3} r_{\alpha\beta} u^\alpha u^\beta - u_\alpha \tau_{\beta\gamma}^\alpha G^{\beta\gamma} \right\}.$$

The last two formulas can be used for a study of effective charges as in [22] where some solutions of RG-equations are analyzed. We shall not consider in this paper such methods connected with the theory of differential equations.

C. Two-loop β -functions for the LAS-model

In order to obtain two-loops of the LAS-model we add to the list (3.9) the expansion

$$\Delta I_{1|2}^{(c)} = -\frac{1}{2} I_1 \int d^2 z \left\{ \hat{r}_{\alpha\beta} \left(\eta^{AB} - \epsilon^{AB} \right) \left(\widehat{\nabla}_A \zeta^\alpha \right) \left(\widehat{\nabla}_B \zeta^\beta \right) + \left(\nabla_\tau \hat{r}_{\alpha\beta} + \hat{r}_{\alpha\gamma} G^{\gamma\delta} T_{[\delta\beta\tau]} \right) \times \right.$$

$$\begin{aligned}
& \left(\eta^{AB} - \epsilon^{AB} \right) \zeta^\tau \left(\widehat{\nabla}_A \zeta^\alpha \right) \partial_B u^\beta + \left(\nabla_\tau \widehat{r}_{\alpha\beta} - T_{[\alpha\tau\gamma]} \widehat{r}_{\beta}^{\gamma} \right) \left(\eta^{AB} - \epsilon^{AB} \right) \partial_A u^\alpha \left(\widehat{\nabla}_B \zeta^\beta \right) \zeta^\tau + \\
& \left(\eta^{AB} - \epsilon^{AB} \right) \left(\frac{1}{2} \nabla_\gamma \nabla_\tau \widehat{r}_{\alpha\beta} + \frac{1}{2} \widehat{r}_{\epsilon\beta} r_{\gamma\tau\alpha}^\epsilon + \frac{1}{2} \widehat{r}_{\alpha\epsilon} r_{\gamma\tau\beta}^\epsilon + T_{[\alpha\tau\epsilon]} \widehat{r}^{\epsilon\delta} T_{[\delta\gamma\beta]} + \right. \\
& \left. G^{\mu\nu} T_{[\nu\beta\gamma]} \nabla_\tau \widehat{r}_{\alpha\mu} - G^{\mu\nu} T_{[\alpha\gamma\nu]} \nabla_\tau \widehat{r}_{\mu\beta} \right) \partial_A u^\alpha \partial_B u^\beta \zeta^\tau \zeta^\gamma \}
\end{aligned}$$

and the d-covariant part of the expansion for the one-loop mass counter-term

$$\Delta I_{1|2}^{(m)} = \frac{(\tilde{m})^2}{2} I_1 \int d^2 z \frac{1}{3} r_{\alpha\beta} \zeta^\alpha \zeta^\beta.$$

The non-distinguished diagrams defining two-loop divergences are illustrated in Fig.2. We present the explicit form of corresponding counter-terms computed by using, in our case adapted to la-backgrounds, methods developed in [27,29]:

For counter-term of the diagram (α) we obtain

$$\begin{aligned}
(\alpha) = & -\frac{1}{2} \lambda^2 (I_i)^2 \int d^2 z \{ \left(\frac{1}{4} \Delta r_{\delta\varphi} - \frac{1}{12} \nabla_\delta \nabla_\varphi \overleftarrow{r} + \frac{1}{2} r_{\delta\alpha} r_\varphi^\alpha - \frac{1}{6} r_{\delta\varphi}^{\alpha\beta} r_{\alpha\beta} + \right. \\
& + \frac{1}{2} r_{\delta\varphi}^{\alpha\beta\gamma} r_{\alpha\varphi\beta\gamma} + \frac{1}{2} G_{\delta\tau} T^{[\tau\alpha\beta]} \Delta T_{[\varphi\alpha\beta]} + \frac{1}{2} G_{\varphi\tau} r_\delta^\alpha T_{[\alpha\beta\gamma]} T^{[\tau\beta\gamma]} - \\
& \frac{1}{6} G^{\beta\tau} T_{[\delta\alpha\beta]} T_{[\varphi\gamma\tau]} r^{\alpha\gamma} + G^{\gamma\tau} T_{[\delta\alpha\tau]} \nabla^{(\alpha} \nabla^{\beta)} T_{[\varphi\beta\gamma]} + \frac{3}{4} G^{\kappa\tau} r_{\delta\varphi}^{\alpha\beta\gamma} T_{[\beta\gamma\kappa]} T_{[\alpha\varphi\tau]} - \\
& \frac{1}{4} r^{\kappa\alpha\beta\gamma} T_{[\delta\beta\gamma]} T_{[\kappa\alpha\varphi]} \} \partial_A u^\delta \partial^A u^\varphi + \frac{1}{4} [\nabla^\beta \Delta T_{[\delta\varphi\beta]} - 3 r_{\dots\delta}^{\gamma\beta\alpha} \nabla_\alpha T_{[\beta\gamma\varphi]} - 3 T_{[\alpha\beta\delta]} \nabla^\gamma r_{\dots\gamma\varphi}^{\beta\alpha} + \\
& \frac{1}{4} r^{\alpha\gamma} \nabla_\alpha T_{[\gamma\delta\varphi]} + \frac{1}{6} T_{[\delta\varphi\alpha]} \nabla^\alpha r - 4 G^{\gamma\tau} T_{[\tau\beta\delta]} T_{[\alpha\kappa\varphi]} \nabla^\beta (G_{\gamma\epsilon} T^{[\alpha\kappa\epsilon]}) + \\
& \left. 2 G_{\delta\tau} G^{\beta\epsilon} \nabla_\alpha (G^{\alpha\nu} T_{[\nu\beta\gamma]}) T^{[\gamma\kappa\tau]} T_{[\epsilon\kappa\varphi]} \right] \epsilon^{AB} \partial_A u^\delta \partial_B u^\varphi \}.
\end{aligned}$$

In order to computer the counter-term for diagram (β) we use integrals:

$$\lim_{u \rightarrow v} i < \partial_A \zeta(u) \partial^A \zeta(v) > = i \int \frac{d^q p}{(2\pi)^q} \frac{p^2}{p^2 - \tilde{m}^2} = \tilde{m}^2 I_1$$

(containing only a IR-divergence) and

$$J \equiv i \int \frac{d^2 p}{(2\pi)^2} \frac{1}{(p^2 - \tilde{m}^2)^2} = -\frac{1}{(2\pi)^2} \int d^2 k_E \frac{1}{(k_E^2 + \tilde{m}^2)^2}.$$

(being convergent). In result we can express

$$(\beta) = \frac{1}{6} \lambda^2 (I_1^2 + 2\tilde{m} I_1 J) \int d^2 z \hat{r}_{\beta\alpha\gamma\delta} r^{\beta\gamma} (\eta^{AB} - \epsilon^{AB}) \partial_A u^\alpha \partial_B u^\beta.$$

In our further considerations we shall use identities (we can verify them by straightforward calculations):

$$\hat{r}_{(\alpha,\beta)}^{[\gamma,\delta]} = -\nabla_{(\alpha}(G_{\beta)\tau} T^{[\tau\gamma\delta]}), \hat{r}_{[\beta\alpha\gamma\delta]} = 2G^{\kappa\tau} T_{[\tau[\alpha\beta]T_{[\gamma\delta]\kappa]},$$

in the last expression we have three type of antisymmetrizations on indices, $[\tau\alpha\beta], [\gamma\delta\kappa]$ and $[\alpha\beta\gamma\delta]$,

$$\begin{aligned} \widehat{\nabla}_\delta T_{[\alpha\beta\gamma]} \widehat{\nabla}_\varphi T^{[\alpha\beta\gamma]} &= \frac{9}{16} (\hat{r}_{[\beta\alpha\gamma]\delta} - \hat{r}_{\delta[\alpha\beta\gamma]}) (\hat{r}_{\dots\dots\dots\varphi}^{[\beta\alpha\gamma]} - \hat{r}_{\dots\dots\dots\varphi}^{[\alpha\beta\gamma]}) - \\ &- \frac{9}{4} \hat{r}_{[\alpha\beta\gamma\delta]} \hat{r}_{\dots\dots\dots\varphi}^{[\alpha\beta\gamma]} + \frac{9}{4} \hat{r}_{\dots\dots\dots[\delta} \hat{r}_{[\varphi]\alpha\beta\gamma]}^{\alpha\beta\gamma} + \frac{9}{4} \hat{r}_{[\delta}^{\alpha,\beta\gamma} \hat{r}_{[\varphi]\alpha\beta\gamma]}. \end{aligned} \quad (4.3)$$

The momentum integral for the first of diagrams (γ)

$$\int \frac{d^q p d^q p'}{(2\pi)^{2q}} \frac{p_A p_B}{(p^2 - \tilde{m}^2)([k+q]^2 - \tilde{m}^2)([p+q]^2 - \tilde{m}^2)}$$

diverges for a vanishing exterior momenta k_μ . The explicit calculus of the corresponding counter-term results in

$$\begin{aligned} \gamma_1 &= -\frac{2\lambda^2}{3q} I_1^2 \int d^2 z \{ (r_{\alpha(\beta\gamma)\delta} + G^{\varphi\tau} T_{[\tau\alpha(\beta]T_{[\gamma]\delta\varphi]})} (r_{\mu}^{\beta,\gamma\delta} - G_{\mu\tau} G_{\kappa\epsilon} T^{[\tau\beta\kappa]} T^{[\gamma\delta\epsilon]}) \partial^A u^\alpha \partial_A u^\mu + \\ &(\nabla_{(\beta} T_{[\delta]\alpha\gamma]) \nabla^\beta (G_{\mu\tau} T^{[\tau\gamma\delta]}) \epsilon^{LN} \eta_{NM} \epsilon^{MR} \partial_L u^\alpha \partial_R u^\mu - \\ &2(r_{\alpha(\beta\gamma)\delta} + G^{\varphi\tau} T_{[\alpha\tau(\beta]T_{[\gamma]\delta\varphi]}) \nabla^\beta (G_{\mu\epsilon} T^{[\epsilon\delta\gamma]}) \epsilon^{MR} \partial_M u^\alpha \partial_R u^\mu \}. \end{aligned} \quad (4.4)$$

The counter-term of the sum of next two (γ)-diagrams is chosen to be the la-extension of that introduced in [29,27]:

$$\gamma_2 + \gamma_3 = -\lambda^2 \omega(\epsilon) \frac{10-7q}{18q} I_1^2 \int d^2 z \{ \widehat{\nabla}_A T_{[\alpha\beta\gamma]} \widehat{\nabla}^A T^{[\alpha\beta\gamma]} +$$

$$6G^{\tau\epsilon}T_{[\delta\alpha\tau]}T_{[\epsilon\beta\gamma]}\widehat{\nabla}_\varphi T^{[\alpha\beta\gamma]}(\eta^{AB}-\epsilon^{AB})\partial_A u^\delta\partial_B u^\varphi\}. \quad (4.5)$$

In a similar manner we can computer the rest part of counter-terms:

$$\begin{aligned} \delta &= \frac{1}{2}\lambda^2(I_1^2 + m^2 I_1 J) \int d^2 z \widehat{r}_{\alpha(\beta\gamma)\delta} \widehat{r}^{\beta\gamma} (\eta^{AB} - \epsilon^{AB}) \partial_A u^\alpha \partial_B u^\delta, \\ \epsilon &= \frac{1}{4}\lambda^2 I_1^2 \int d^2 z (\eta^{AB} - \epsilon^{AB}) [\Delta \widehat{r}_{\delta\varphi} + r_\delta^\alpha \widehat{r}_{\alpha\varphi} + r_\varphi^\alpha \widehat{r}_{\delta\alpha} \\ &\quad - 2(G^{\alpha\tau} T_{[\delta\beta\alpha]} T_{[\varphi\gamma\tau]} \widehat{r}^{\beta\gamma} - G_{\varphi\tau} T^{[\tau\alpha\beta]} \nabla_\alpha \widehat{r}_{\delta\beta} + G_{\delta\tau} T^{[\tau\alpha\beta]} \nabla_\alpha \widehat{r}_{\beta\varphi}] \partial_A u^\delta \partial_B u^\varphi, \\ \iota &= \frac{1}{6}\lambda^2 \tilde{m}^2 I_1 J \int d^2 z \widehat{r}_{\beta\alpha\gamma\delta} r^{\beta\gamma} (\eta^{AB} - \epsilon^{AB}) \partial_A u^\alpha \partial_B u^\delta, \\ \eta &= \frac{1}{4}\lambda^2 \omega(\epsilon) (I_1^2 + 2\tilde{m}^2 I_1 J) \int d^2 z \widehat{r}_{\alpha\gamma\delta}^\beta T_{[\beta\varphi\tau]} T^{[\gamma\varphi\tau]} (\eta^{AB} - \epsilon^{AB}) \partial_A u^\alpha \partial_B u^\delta. \end{aligned}$$

By using relations (4.3) we can represent terms (4.4) and (4.5) in the canonical form (4.1) from which we find the contributions in the $\beta_{\delta\varphi}$ -function (4.2):

$$\begin{aligned} \gamma_1 &: -\frac{2}{3(2\pi)^2} \widehat{r}_{\delta(\alpha\beta)\gamma} \widehat{r}^{\gamma(\alpha\beta)}_{\dots\dots\dots\varphi} - \frac{(\omega_1 - 1)}{(2\pi)^2} \left\{ \frac{4}{3} \widehat{r}_{[\gamma(\alpha\beta)\delta]} \widehat{r}^{[\gamma(\alpha\beta)]}_{\dots\dots\dots\varphi} + \widehat{r}_{[\alpha\beta\gamma\delta]} \widehat{r}^{[\alpha\beta\gamma]}_{\dots\dots\dots\varphi} \right\}, \\ \gamma_2 + \gamma_3 &: \frac{(4\omega_1 - 5)}{9(2\pi)^2} \left\{ \widehat{\nabla}_\delta T_{[\alpha\beta\gamma]} \widehat{\nabla}_\varphi T^{[\alpha\beta\gamma]} + 6G^{\tau\epsilon} T_{[\delta\alpha\tau]} T_{[\epsilon\beta\gamma]} \widehat{\nabla}_\varphi T^{[\alpha\beta\gamma]} - \frac{(\omega_1 - 1)}{(2\pi)^2} \widehat{r}_{[\alpha\beta\gamma\delta]} \widehat{r}^{[\alpha\beta\gamma]}_{\dots\dots\dots\varphi} \right\}, \end{aligned} \quad (4.8)$$

$$\eta : \frac{\omega_1}{(2\pi)^2} \widehat{r}^{\alpha}_{\cdot\delta\beta\varphi} T_{[\alpha\tau\epsilon]} T^{[\beta\tau\epsilon]}.$$

Finally, in this subsection we remark that two-loop β -function can not be written only in terms of curvature $\widehat{r}_{\alpha\beta\gamma\delta}$ and d-derivation $\widehat{\nabla}_\alpha$ (similarly as in the locally isotropic case [29,27]).

D. Low-Energy Effective Action for La-Strings

The conditions of vanishing of β -functions describe the propagation of string in the background of la-fields $G_{\alpha\beta}$ and $b_{\alpha\beta}$. (in this section we chose the canonic d-connection ${}^\circ\Gamma_{\beta\gamma}^\alpha$ on $\mathcal{E}$). The β -functions are proportional to d-field equations obtained from the on-shell string effective action

$$I_{eff} = \int du \sqrt{|\gamma|} L_{eff}(\gamma, b). \quad (4.9)$$

The adapted to N-connection variations of (4.8) with respect to $G^{\mu\nu}$ and $b^{\mu\nu}$ can be written as

$$\begin{aligned} \frac{\Delta I_{eff}}{\delta G^{\alpha\beta}} &= W_{\alpha\beta} + \frac{1}{2} G_{\alpha\beta} (L_{eff} + \text{complete derivation}), \\ \frac{\Delta I_{eff}}{\delta b^{\alpha\beta}} &= 0. \end{aligned}$$

The invariance of action (4.8) with respect to N-adapted diffeomorfisms gives rise to the identity

$$\begin{aligned} \nabla_\beta W^{\alpha\beta} - T^{[\alpha\beta\gamma]} \frac{\Delta I_{eff}}{\delta b^{\beta\gamma}} = \\ -\frac{1}{2} \nabla^\alpha (L_{eff} + \text{complete derivation}) \end{aligned}$$

(in the locally isotropic limit we obtain the well-known results from [33,34]). This points to the possibility to write out the integrability conditions as

$$\nabla^\beta \beta_{(\alpha\beta)} - G_{\alpha\tau} T^{[\tau\beta\gamma]} \beta_{[\beta\gamma]} = -\frac{1}{2} \nabla_\alpha L_{eff}. \quad (4.10)$$

For one-loop β -function, $\beta_{\alpha\beta}^{(1)} = \frac{1}{2\pi} \hat{r}_{\alpha\beta}$, we find from the last equations

$$\nabla^\beta \beta_{(\delta\beta)}^{(1)} - G_{\delta\tau} T^{[\tau\beta\gamma]} \beta_{[\beta\gamma]}^{(1)} = \frac{1}{4\pi} \nabla_\delta \left(\overleftarrow{R} + \frac{1}{3} T_{[\alpha\beta\gamma]} T^{[\alpha\beta\gamma]} \right)$$

We can take into account two-loop β -functions by fixing an explicit form of

$$\omega(\epsilon) = 1 + 2\omega_1\epsilon + 4\omega_2\epsilon^2 + \dots$$

when

$$\omega_{HVB}(\epsilon) = \frac{1}{(1-\epsilon)^2}, \omega_1^{HVB} = 1, \omega_2^{HVB} = \frac{3}{4}$$

(the t'Hooft-Veltman-Bos prescription [35]). Putting values (4.8) into (4.10) we obtain the two-loop approximation for la-field equations

$$\begin{aligned} \nabla^\beta \beta_{(\delta\beta)}^{(2)} - G_{\delta\tau} T^{[\tau\beta\gamma]} \beta_{[\beta\gamma]}^{(2)} &= \frac{1}{2(2\pi)^2} \nabla_\delta \left[-\frac{1}{8} r_{\alpha\beta\gamma\delta} r^{\alpha\beta\gamma\delta} + \frac{1}{4} r_{\alpha\beta\gamma\delta} G_{\tau\epsilon} T^{[\alpha\beta\tau]} T^{[\epsilon\gamma\delta]} + \right. \\ &\quad \left. \frac{1}{4} G^{\beta\epsilon} G_{\alpha\kappa} T^{[\alpha\tau\sigma]} T_{[\epsilon\tau\sigma]} T^{[\kappa\mu\nu]} T_{[\beta\mu\nu]} - \frac{1}{12} G^{\beta\epsilon} G_{\gamma\lambda} T_{[\alpha\beta\tau]} T^{[\alpha\gamma\varphi]} T_{[\epsilon\varphi\kappa]} T^{[\lambda\tau\kappa]} \right], \end{aligned}$$

which can be obtained from effective action

$$\begin{aligned} I^{eff} \sim \int \delta^{n+m} u \sqrt{|\gamma|} \left[-\frac{1}{3} T_{[\alpha\beta\gamma]} T^{[\alpha\beta\gamma]} - \frac{\alpha'}{4} \left(\frac{1}{2} r_{\alpha\beta\gamma\delta} r^{\alpha\beta\gamma\delta} - \right. \right. \\ \left. \left. G_{\tau\epsilon} r_{\alpha\beta\gamma\delta} T^{[\alpha\beta\tau]} T^{[\epsilon\gamma\delta]} - G^{\beta\epsilon} G_{\alpha\nu} T^{[\alpha\gamma\kappa]} T_{[\epsilon\gamma\kappa]} T^{[\nu\sigma\varsigma]} T_{[\beta\sigma\varsigma]} + \right. \right. \\ \left. \left. \frac{1}{3} G^{\beta\epsilon} G_{\delta\kappa} T_{[\alpha\beta\gamma]} T^{[\alpha\delta\varphi]} T_{[\varphi\nu\epsilon]} T^{[\gamma\nu\kappa]} \right) \right]. \end{aligned} \quad (4.11)$$

The action (4.11) (for $2\pi\alpha' = 1$ and in locally isotropic limit) is in good concordance with the similar ones on usual closed strings [36,27].

We note that the existence of an effective action is assured by the Zamolodchikov c-theorem [37] which was generalized [38] for the case of bosonic nonlinear σ -model with dilaton connection. In a similar manner we can prove that such results hold good for la-backgrounds.

V. SCATTERING OF GRAVITONS AND DUALITY OF LOCALLY ANISOTROPIC SIGMA-MODELS

The quantum theory of la-strings can be naturally considered by using the formalism of functional integrals on "hypersurfaces" (see Polyakov's works [39]). In this section we study the structure of scattering amplitudes of la-gravitons. Questions on duality of la-string theories will be also analyzed.

A. The generation functional of la-string amplitudes for scattering of la-gravitons

We introduce the Green function of la-tachyons, the fundamental state of la-string, as an integral (after Weeck rotation in the Euclidean space)

$$G_t(p_1, \dots, p_N) = \int [D\gamma_{AB}(\zeta)][Du^\alpha(z)] \exp\left(-\frac{1}{4\pi\alpha'} \int d^2z \sqrt{|\gamma|} \gamma^{AB} \partial_A u^\alpha \partial_B u^\alpha\right) \int \left[\prod_A D^2 z_A \right] \sqrt{|\gamma(z_B)|} \exp(ip_B^\alpha u(z_B)), \quad (5.1)$$

where the integration measure on γ_{AB} includes the standard ghost Fadeev-Popov determinant corresponding to the fixation of the reparametrization invariance and

$$\left[\prod_A D^2 z_A \right] = \prod_{A \neq M, N, K} d^2 z_A |z_M - z_N|^2 |z_K - z_M|^2 |z_K - z_N|^2.$$

Because the string quantum field theory can be uncontradictory formulated for spaces of dimension $d=26$ we consider that in formula (5.1) α takes values from 0 to 25. Formula (5.1) leads to dual amplitudes for la-tachyon scatterings for $p^2 = \frac{4}{\alpha'}$ (see [40] for details and references on usual locally isotropic tachyon scattering).

The generating functional of Green functions (5.1) in the coordinate u -representation can be formally written as a hyper surface mean value

$$\Gamma^0[\Phi] = \langle \exp \left(-\frac{1}{2\pi\alpha'} \int d^2z \sqrt{|\gamma|} \Phi[u(z)] \right) \rangle .$$

In order to conserve the reparametrization invariance we define the la-graviton source as

$$\Gamma^0[G] = \langle \exp \left(-\frac{1}{4\pi\alpha'} \int d^2z \sqrt{|\gamma|} \gamma^{AB} \partial_A u^\alpha \partial_B u^\beta G_{\alpha\beta}[u(z)] \right) \rangle$$

from which we obtain the Green function of a number of K elementary perturbations of the closed la-string (K la-gravitons)

$$G_g(u_1, \dots, u_K) = \langle \frac{1}{2} \int \left[\prod_{[j]} D^2 z_{[j]} \right] \sqrt{|\gamma(z)|} G^{AB}(z_{[j]}) \partial_A u^\alpha(z_{[j]}) \partial_B u^\beta(z_{[j]}) \chi_{\alpha\beta}^{(j)} \delta^{(d)}(u_{[j]} - u(z_{[j]})) \rangle ,$$

where $\chi_{\alpha\beta}^{(j)}$ are polarization d-tensors of exterior la-gravitons and $[j] = 1, 2, \dots, K$. Applying the Fourier transform we obtain

$$G_g(p_1, \dots, p_K) = \int [D\gamma_{AB}(z)] [Du^\alpha(z)] \exp \left(-\frac{1}{2\pi\alpha'} \int d^2z \sqrt{|\gamma|} \frac{1}{2} G^{AB} \partial_A u^\alpha \partial_B u^\alpha \right) \\ \int \left[\prod_{[j]} D^2 z_{[j]} \right] \frac{1}{2} \sqrt{|\gamma(z_{[j]})|} \gamma^{AB}(z_{[j]}) \partial_A u^\beta(z_{[j]}) \partial_B u^\gamma(z_{[j]}) \chi_{\alpha\beta}^{(j)} \exp[ip_{[j]}^\delta u^\delta(z)] .$$

Integrating the last expression on $G_{\mu\nu}$ and u^α for $d=26$, when there are not conformal anomalies, we have

$$G_g(p_1, \dots, p_K) = \int \left[\prod_{[j]} D^2 z_{[j]} \right] \left(\partial_A \frac{\partial}{\partial p_{[j]}} \cdot \chi^{(j)} \cdot \partial^A \frac{\partial}{\partial p_{[j]}} \right) \times \\ \int [D\sigma(z)] \exp[-\pi\alpha' \sum_{[i,j]} p_{[i]} p_{[j]} V(z_{[i]}, z_{[j]}, \sigma)] , \quad (5.2)$$

where V is the Green function of the Laplacian for the conformally-flat metric $G_{AB} = e^\sigma \delta_{AB}$:

$$\partial_A (\sqrt{|\gamma|} \gamma^{AB} \partial_A) V = -\delta^2(z_{[i]} - z_{[j]})$$

which can be represented as

$$V(z_{[i]}, z_{[j]}, \sigma) = -\frac{1}{4\pi} \ln |z_{[i]} - z_{[j]}|^2, \quad z_{[i]} \neq z_{[j]},$$

$$V\left(z_{[k]}, z_{[k]}^p, \sigma\right) = \frac{1}{4\pi} \left(\sigma(z_{[k]}) - \ln\left(\frac{1}{\epsilon}\right)\right),$$

for ϵ being the cutting parameter. Putting the last expression into (5.2) we compute the Green function of la-gravitons:

$$G_g(p_1, \dots, p_K) = \int \prod_{[j] \neq [p], [q], [s]} d^2 z_{[j]} |z_{[q]} - z_{[p]}|^2 |z_{[s]} - z_{[q]}|^2 |z_{[s]} - z_{[p]}|^2 \left(\partial_A \frac{\partial}{\partial p_{[j]}} \cdot \chi^{(j)} \cdot \partial^A \frac{\partial}{\partial p_{[j]}} \right)$$

$$\int d\sigma(z) \prod_{[i] < [m]} |z_{[i]} - z_{[m]}|^{\alpha' p_{[i]} p_{[m]}} \exp \left[-\frac{1}{4} \alpha' \sum_{[k]} p_{[k]}^2 \sigma(z_{[k]}) \right] \exp \left[\frac{\alpha'}{4} \ln\left(\frac{1}{\epsilon}\right) \sum_{[k]} p_{[k]}^2 \right],$$

where the definition of integration on $\sigma(z)$ is extended as

$$\int d\sigma(z) \equiv \lim_{\sigma_d(z_{[j]}) \rightarrow -\infty}^{\sigma_s(z_{[j]}) \rightarrow +\infty} \prod_{[j]} \int_{\sigma_d(z_{[j]})}^{\sigma_s(z_{[j]})} d\sigma(z_{[j]}).$$

So the scattering amplitude

$$A_g(p_1, \dots, p_K) = \lim_{p_{[j]}^2 \rightarrow 0} \prod_{[j]} p_{[j]}^2 G_g(p_1, \dots, p_K)$$

is finite if

$$\lim_{\sigma_d(z_{[j]}) \rightarrow -\infty, p_{[j]}^2 \rightarrow 0} |p_{[j]}^2 \sigma(z_{[j]})| = \text{const} < \infty.$$

$G_g(p_1, \dots, p_K)$ has poles on exterior momenta corresponding to massless locally anisotropic modes of spin 2 (la-gravitons). The final result for the scattering amplitude of la-gravitons is of the form

$$A_g(p_1, \dots, p_K) \sim \int \prod_{[j] \neq [p], [q], [s]} d^2 z_{[j]} |z_{[p]} - z_{[q]}|^2 |z_{[s]} - z_{[q]}|^2 |z_{[s]} - z_{[p]}|^2$$

$$\partial_A \frac{\partial}{\partial p_{[j]}^\alpha} \chi_{\alpha\beta}^{(j)} \partial^A \frac{\partial}{\partial p_{[j]}^\beta} \prod_{[m] < [n]} |z_{[m]} - z_{[n]}|^{\alpha' p_{[m]} \cdot p_{[n]}}.$$

If instead of polarization d-tensor $\chi_{\alpha\beta}^{(j)}$ the graviton polarization tensor $\chi_{ik}^{(j)}$ is taken we obtain the well known results on scattering of gravitons in the framework of the first quantization of the string theory [2,41].

B. Duality of La- σ -models

Two theories are dual if their non-equivalent second order actions can be generated by the same first order action. The action principle assures the equivalence of the classical dual theories. But, in general, the duality transforms affects the quantum conformal properties [42]. In this subsection we shall prove this for the la- σ -model (3.1) when metric γ and the torsion potential b on la-background ξ do not depend on coordinate u^0 . If such conditions are satisfied we can write for (3.1) the first order action

$$I = \frac{1}{4\pi\alpha'} \int d^2z \left\{ \sum_{\alpha,\beta=1}^{n+m-1} [\sqrt{|\gamma|} \gamma^{AB} (G_{00} V_A V_B + 2G_{0\alpha} V_A (\partial_B u^\alpha) + G_{\alpha\beta} (\partial_A u^\alpha) (\partial_B u^\beta)) + \right. \\ \left. \epsilon^{AB} (b_{0\alpha} V_B (\partial_A u^\alpha) + b_{\alpha\beta} (\partial_A u^\alpha) (\partial_B u^\beta)) \right] + \epsilon^{AB} \hat{u}^0 (\partial_A V_B) + \alpha' \sqrt{|\gamma|} R^{(2)} \Phi(u) \right\}, \quad (5.3)$$

where string interaction constants from (3.1) and (5.3) are related as $\lambda^2 = 2\pi\alpha'$.

This action will generate an action of type (3.1) if we shall exclude the Lagrange multiplier $\hat{u}^0$. The dual to (5.3) action can be constructed by substituting V_A expressed from the motion equations for fields V_A (also obtained from action (5.3)):

$$\hat{I} = \frac{1}{4\pi\alpha'} \int d^2z \left\{ \sqrt{|\gamma|} \gamma^{AB} \hat{G}_{\alpha\beta} (\partial_A \hat{u}^\alpha) (\partial_B \hat{u}^\beta) + \epsilon^{AB} \hat{b}_{\alpha\beta} (\partial_A \hat{u}^\alpha) (\partial_B \hat{u}^\beta) + \alpha' \sqrt{|\gamma|} R^{(2)} \Phi(u) \right\},$$

where the new metric and torsion potential are introduced respectively as

$$\hat{G}_{00} = \frac{1}{G_{00}}, \hat{G}_{0\alpha} = \frac{b_{0\alpha}}{G_{00}}, \hat{G}_{\alpha\beta} = G_{\alpha\beta} - \frac{G_{0\alpha} G_{0\beta} - b_{0\alpha} b_{0\beta}}{G_{00}}$$

and

$$\hat{b}_{0\alpha} = -\hat{b}_{\alpha 0} = \frac{G_{0\alpha}}{G_{00}}, \hat{b}_{\alpha\beta} = b_{\alpha\beta} + \frac{G_{0\alpha}b_{0\beta} - b_{0\alpha}G_{0\beta}}{G_{00}}$$

(in the formulas for the new metric and torsion potential indices α and β take values $1, 2, \dots, n+m-1$).

If the model (3.1) satisfies the conditions of one-loop conformal invariance (see details for locally isotropic backgrounds in [3]), one holds these la-field equations

$$\begin{aligned} \frac{1}{\alpha'} \frac{n+m-25}{3} + [4(\nabla\Phi)^2 - 4\nabla^2\Phi - r - \frac{1}{3}T_{[\alpha\beta\gamma]}T^{[\alpha\beta\gamma]}] &= 0, \\ \hat{r}_{(\alpha\beta)} + 2\nabla_{(\alpha}\nabla_{\beta)}\Phi &= 0, \\ \hat{r}_{[\alpha\beta]} + 2T_{[\alpha\beta\gamma]}\nabla^\gamma\Phi &= 0. \end{aligned} \tag{5.4}$$

By straightforward calculations we can show that the dual theory has the same conformal properties and satisfies the conditions (5.4) if the dual transform is completed by the shift of dilaton field

$$\hat{\Phi} = \Phi - \frac{1}{2} \log G_{00}.$$

The system of la-field equations (5.4), obtained as a low-energy limit of the la-string theory, is similar to Einstein-Cartan equations (2.28) and (2.29). We note that the explicit form of locally anisotropic energy-momentum source in (5.4) is defined from well defined principles and symmetries of string interactions and this form is not postulated, as in usual locally isotropic field models, from some general considerations in order to satisfy the necessary conservation laws on la-space whose formulation is very sophisticated because of nonexistence of global and even local group of symmetries of such type of spaces. Here we also remark that the LAS-model with dilaton field interactions does not generate in the low-energy limit the Einstein-Cartan la-theory because the first system of equations from

(5.4) represents some constraints (being a consequence of the two-dimensional symmetry of the model) on torsion and scalar curvature which can not be interpreted as some algebraic relations of type (2.29) between locally anisotropic spin-matter source and torsion. As a matter of principle we can generalize our constructions by introducing interactions with gauge la-fields and considering a variant of locally anisotropic chiral σ -model [43] in order to get a system of equations quite similar to (2.29). However, there are not exhaustive arguments for favoring the Einstein-Cartan theory and we shall not try in this work to generate it necessarily from la-strings.

VI. SUMMARY AND CONCLUSIONS

Let us try to summarize our results, discuss their possible implications and make the basic conclusions. Firstly, we have shown that the Einstein-Cartan theory has a natural extension for a various class of la-spaces. Following the R. Miron and M. Anastesiei approach [10] to the geometry of la-spaces it becomes evident the possibility and manner of formulation of classical and quantum field theories on such spaces. Here we note that in la-theories we have an additional geometric structure, the N-connection. From physical point of view it can be interpreted, for instance, as a fundamental field managing the dynamics of splitting of high-dimensional space-time into the four-dimensional and compactified ones. We can also consider the N-connection as a generalized type of gauge field which reflects some specifics of la-field interactions and possible intrinsic structure of la-spaces. It was convenient to analyze the geometric structure of different variants of la-spaces (for instance, Finsler, Lagrange and generalized Lagrange spaces) in order to make obvious physical properties and compare theirs perspectives in developing of new physical models.

According to modern-day views the theories of fundamental field interactions should be a low energy limit of the string theory. One of the main results of this work is the proof of the fact that in the framework of la-string theory is contained a more general, locally anisotropic, gravitational physics. To do this we have developed the locally anisotropic nonlinear sigma model and studied its general properties. We shown that the condition of self consistent propagation of string on la-background impose corresponding constraints on the N-connection curvature, la-space torsion and antisymmetric d-tensor. Our extension of background field method for la-spaces has a distinguished by N-connection character and the main advantage of this formalism is doubtlessly its universality for all types of locally isotropic or anisotropic spaces.

The presented one- and two-loop calculus for the LAS-model and used in this work d-covariant dimensional regularization are developed for la-background spaces modelled as vector bundles provided with compatible N-connection, d-connection and metric structures. In the locally isotropic limit we obtain the corresponding formulas for the usual nonlinear sigma model.

Finally, it should be stressed that we firstly calculated the amplitudes for scattering of la-gravitons and that the duality properties of the formulated in this work LAS-model are similar to those of models considered for locally isotropic strings

ACKNOWLEDGMENTS

I would like to thank Sergiu V. Ostaf for useful comments and discussions.

REFERENCES

- [1] C. Lovelace, Phys. Lett. B **35**, 75 (1984).
- [2] S. Fradkin and A. A. Tseytlin, Phys. Lett. B **158**, 316 (1985); Nucl. Phys. B **261**, 1 (1985).
- [3] C. G. Callan, D. Friedan, E. J. Martinec, and M. J. Perry, Nucl. Phys. B **262**, 593 (1985).
- [4] A. Sen, Phys. Rev. Lett. **55**, 1846 (1986).
- [5] S. P. de Alwis, Phys. Rev. D **34**, 3760 (1986).
- [6] A. M. Polyakov, Phys. Lett. B **103**, 107 (1981).
- [7] P. Finsler, *Über und Flächen in allgemeinen Räumen*. Dissertation, Göttingen, 1918 (Verlag Birkhäuser, Basel, 1951).
- [8] L. Berwalld, Math. Z. **25**, 40 (1926).
- [9] E. Cartan, *Les Espaces de Finsler*, (Hermann, Paris, 1934).
- [10] R. Miron and M. Anastasiei, *The Geometry of Lagrange Spaces: Theory and Applications* (Kluwer, Dordrecht, 1994); R. Miron and M. Anastasiei, *Vector Bundles. Lagrange Spaces. Applications to Relativity* (Editura Academiei Române, Bucharest, 1987) (in Romanian).
- [11] H. Rund, *The Differential Geometry of Finsler Spaces*, (Springer-Verlag, Berlin, 1959); G. S. Asanov, *Finsler Geometry, Relativity and Gauge Theories*, (D. Reidel, Dordrecht, 1985); G. S. Asanov and S. F. Ponomarenko *Finsler Bundles on Space-Time. Associated*

- Gauge Fields and Connections*, (Știința, Chișinău, 1988) (in Russian); R. Miron and T. Kawaguchi, Int. J. Theor. Phys. **30**, 1521 (1991); A. A. Vlasov, *Statistical Distribution Functions* (Nauka, Moscow, 1966) (in Russian); R. Pimenov, Spaces of Kinematic Type (Nauka, Leningrad, 1968) (in Russian); G. Yu. Bogoslovskiy, The Theory of Locally Anisotropic Space-Time, (Moscow University Press, Moscow, 1982) (in Russian); A. Bejancu, *Finsler Geometry and Applications* (Ellis Horwood Chichester, England, 1990).
- [12] S. Vacaru, J. Math. Phys. **37**, 508 (1996).
- [13] S. Vacaru and Yu. Goncharenko, Int. J. Theor. Phys. **34**, 1955 (1995).
- [14] S. Vacaru, Nonlinear Connections in Superbundles and Locally Anisotropic Supergravity (in preparation).
- [15] J. Kern, Arch. Math. **25**, 438 (1974).
- [16] W. Barthel, J. Reine. Angew. Math. **212**, 120 (1963).
- [17] M. Anastasiei, An. Șt. Univ. Iași, s I a **32**, 17 (1986).
- [18] A. A. Tseytlin, Int. J. Mod. Phys. A **4**, 1257 (1989).
- [19] C. M. Hull, *Lectures on Non-linear Sigma Models and Strings* (DAMPT, Cambridge, England, 1987).
- [20] E. Witten, Commun. Math. Phys. **92**, 455 (1984).
- [21] J. Wess and B. Zumino, Phys. Lett. **37**, 95 (1971).
- [22] E. Braaten, T. L. Curtright, C. K., and Zachos, Nucl. Phys. B **260**, 630 (1985).

- [23] A. A. Belavin, A. M. Polyakov, and A. B. Zamolodchikov, Nucl. Phys. B **241**, 333 (1984).
- [24] S. V. Ketov, *Introduction into Quantum Theory of Strings and Superstrings* (Nauka, Novosibirsk, 1990) (in Russian).
- [25] B. S. DeWitt, *Dynamical Theory of Groups and Fields* (Gordon and Breach, London, 1965).
- [26] L. Alvarez-Gaume, D. Z. Freedman, and S. Mukhi, Ann. Phys. **134**, 85 (1981).
- [27] S. V. Ketov, *Nonlinear Sigma-Models in Quantum Field Theory and String Theory* (Nauka, Novosibirsk, 1992) (in Russian).
- [28] R. S. Howe, and K. S. Stelle, The Background Field Method and the Nonlinear σ -model, Preprint IAS, 1986 (unpublished).
- [29] S. V. Ketov, Nucl. Phys. **294**, 813 (1987).
- [30] B. E. Fridling and E. A. M. Van de Ven, Nucl. Phys. B **268**, 719 (1986).
- [31] G. t'Hooft, Nucl. Phys. B **61**, 455 (1973).
- [32] T. L. Curtright and C. K. Zachos, Phys. Rev. Lett. **53**, 1799 (1984).
- [33] D. Zanon, Phys. Lett. B **191**, 363 (1987).
- [34] C. G. Callan, I. R. Klebanov, and M. J. Perry, Nucl. Phys. B **278**, 78 (1986).
- [35] G. t'Hooft and M. Veltman, Nucl. Phys. B **44**, 189 (1971); M. Bos, Ann. Phys. (USA) **181**, 177 (1988).

- [36] D. J. Gross and J. H. Sloan, Nucl. Phys. B **291**, 41 (1987).
- [37] A. B. Zamolodchikov, Pis'ma J. T. P. **43**, 565 (1986).
- [38] A. A. Tseytlin, Phys. Lett. B **194**, 63 (1987).
- [39] A. M. Polyakov, Phys. Lett. B **103**, 207 and 211 (1981); *Gauge Fields and Strings*, Contemporary Concepts in Physics (Harwood Acad. Publ., Chur e. a., Switzerland, 1987).
- [40] *Superstrings: the first 15 years of superstring theory: reprints & commentary* by J. H. Schwarz (World Sci., Singapore e. a. 1985).
- [41] S. V. Ketov, Izvestya Vuzov. Fizika **31**, 17 (1988) (in Russian).
- [42] T. H. Buscher, Phys. Lett. B **201**, 466 (1988).
- [43] I. Jack, Nucl. Phys. B **234**, 365 (1984).

FIGURES




FIG. 1. The Feynman diagrams for the one-loop β -functions of the LAS-model




Diagram (α)





Diagram (β)



Diagrams (γ)




Diagram (ε)




Diagram (δ)




Diagram (ι)




Diagram (η)

FIG. 2. Two-loops diagrams for the LAS-model